



# NLK Monoclonal Antibody

|                           |  |
|---------------------------|--|
| <b>Catalog No</b>         | YP-mAb-06124   |
| <b>Isotype</b>            | IgG  |
| <b>Reactivity</b>         | Human;Rat;Mouse  |
| <b>Applications</b>       | WB   |
| <b>Gene Name</b>          | NLK LAK1   |
| <b>Protein Name</b>       | Serine/threonine-protein kinase NLK (EC 2.7.11.24) (Nemo-like kinase) (Protein LAK1)   |
| <b>Immunogen</b>          | Synthesized peptide derived from human protein . at AA range: 230-310  |
| <b>Specificity</b>        | NLK Monoclonal Antibody detects endogenous levels of protein.  |
| <b>Formulation</b>        | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.   |
| <b>Source</b>             | Monoclonal, Mouse,IgG  |
| <b>Purification</b>       | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   |
| <b>Dilution</b>           | WB 1:500-1:2000  |
| <b>Concentration</b>      | 1 mg/ml  |
| <b>Purity</b>             | ≥90%   |
| <b>Storage Stability</b>  | -20°C/1 year   |
| <b>Synonyms</b>           |  |
| <b>Observed Band</b>      | 57kD   |
| <b>Cell Pathway</b>       | Nucleus . Cytoplasm . Predominantly nuclear. A smaller fraction is cytoplasmic (By similarity). .  |
| <b>Tissue Specificity</b> | Amygdala,Placenta,T-cell,Uterus,   |
| <b>Function</b>           | catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,enzyme regulation:Activated by tyrosine and threonine phosphorylation (By similarity). Activated by activin.,function:Role in cell fate determination, required for differentiation of bone marrow stromal cells. Acts downstream of MAP3K7 and HIPK2 to negatively regulate the canonical Wnt/beta-catenin signaling pathway and the phosphorylation and destruction of the MYB transcription factor. May suppress a wide range of transcription factors by phosphorylation of the coactivator, CREBBP (By similarity). Involved in TGFbeta-mediated mesoderm induction, acting downstream of MAP3K7/TAK1 to phosphorylate STAT3.,PTM:Dually phosphorylated on Thr-291 and Tyr-293, which activates the enzyme.,similarity:Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. MAP kinase subfamily.,similarity:Contain |
| <b>Background</b>         | catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,enzyme regulation:Activated by tyrosine and threonine phosphorylation (By similarity). Activated by activin.,function:Role in   |



cell fate determination, required for differentiation of bone marrow stromal cells. Acts downstream of MAP3K7 and HIPK2 to negatively regulate the canonical Wnt/beta-catenin signaling pathway and the phosphorylation and destruction of the MYB transcription factor. May suppress a wide range of transcription factors by phosphorylation of the coactivator, CREBBP (By similarity). Involved in TGFbeta-mediated mesoderm induction, acting downstream of MAP3K7/TAK1 to phosphorylate STAT3.,PTM:Dually phosphorylated on Thr-291 and Tyr-293, which activates the enzyme.,similarity:Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. MAP kinase subfamily.,similarity:Contains 1 protein kinase domain.,subcellular location:Predominantly nuclear. A smaller fraction is cytoplasmic.,subunit:Interacts with STAT3 (By similarity). Interacts with RNF138/NARF and TCF7L2/TCF4. Interacts with HIPK2 and MYB.,

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**