



SCNAA Monoclonal Antibody

Catalog No	YP-mAb-06200
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB
Gene Name	SCN10A
Protein Name	Sodium channel protein type 10 subunit alpha (Peripheral nerve sodium channel 3) (PN3) (hPN3) (Sodium channel protein type X subunit alpha) (Voltage-gated sodium channel subunit alpha Nav1.8)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	SCNAA Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	215kD
Cell Pathway	Cell membrane ; Multi-pass membrane protein . It can be translocated to the cell membrane through association with S100A10. .
Tissue Specificity	Expressed in the dorsal root ganglia and sciatic nerve.
Function	domain:The sequence contains 4 internal repeats, each with 5 hydrophobic segments (S1,S2,S3,S5,S6) and one positively charged segment (S4). Segments S4 are probably the voltage-sensors and are characterized by a series of positively charged amino acids at every third position.,function:This protein mediates the voltage-dependent sodium ion permeability of excitable membranes. Assuming opened or closed conformations in response to the voltage difference across the membrane, the protein forms a sodium-selective channel through which sodium ions may pass in accordance with their electrochemical gradient. It is a tetrodotoxin-resistant sodium channel isoform. Its electrophysiological properties vary depending on the type of the associated beta subunits (in vitro). Plays a role in neuropathic pain mechanisms.,PTM:Ubiquitinated by NEDD4L; which promotes its endocytosis.,similarity:Belongs to t
Background	The protein encoded by this gene is a tetrodotoxin-resistant voltage-gated sodium channel alpha subunit. The properties of the channel formed by the



encoded transmembrane protein can be altered by interaction with different beta subunits. This protein may be involved in the onset of pain associated with peripheral neuropathy. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jun 2014],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images