



# NAC3 Monoclonal Antibody

Catalog No	YP-mAb-06205
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	SLC8A3 NCX3
Protein Name	Sodium/calcium exchanger 3 (Na <sup>+</sup> )/Ca <sup>2+</sup> -exchange protein 3)
Immunogen	Synthesized peptide derived from human protein . at AA range: 530-610
Specificity	NAC3 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	101kD
Cell Pathway	Cell membrane ; Multi-pass membrane protein . Perikaryon . Cell projection, dendrite . Cell projection, dendritic spine . Cell membrane, sarcolemma . Cytoplasm, sarcoplasm . Cell junction . Mitochondrion outer membrane ; Multi-pass membrane protein . Cytoplasm, perinuclear region . Endoplasmic reticulum membrane ; Multi-pass membrane protein . Detected at neuromuscular junctions. .
Tissue Specificity	Isoform 2 is expressed in brain and skeletal muscle. Isoform 3 is expressed in excitable cells of brain, retina and skeletal muscle. Isoform 4 is expressed in skeletal muscle.
Function	enzyme regulation:By intracellular calcium ions.,function:Rapidly transports Ca <sup>2+</sup> during excitation-contraction coupling. Ca <sup>2+</sup> is extruded from the cell during relaxation so as to prevent overloading of intracellular stores.,similarity:Belongs to the sodium/potassium/calcium exchanger family. SLC8 subfamily.,similarity:Contains 2 Calx-beta domains.,tissue specificity:Isoform 2 is expressed in brain and skeletal muscle; Isoform 3 is expressed in excitable cells of brain, retina and skeletal muscle; Isoform 4 is expressed in skeletal muscle.,

**Background**

This gene encodes a member of the sodium/calcium exchanger integral membrane protein family. Na<sup>+</sup>/Ca<sup>2+</sup> exchange proteins are involved in maintaining Ca<sup>2+</sup> homeostasis in a wide variety of cell types. The protein is regulated by intracellular calcium ions and is found in both the plasma membrane and intracellular organellar membranes, where exchange of Na<sup>+</sup> for Ca<sup>2+</sup> occurs in an electrogenic manner. Alternative splicing has been observed for this gene and multiple variants have been described. [provided by RefSeq, Aug 2013],

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**