





SC5A5 Monoclonal Antibody

Catalog No Isotype Reactivity Applications Gene Name Protein Name Immunogen Specificity Formulation Source	IgG Human;Rat;Mouse; WB SLC5A5 NIS Sodium/iodide cotransporter (Na(+)/I(-) cotransporter) (Sodium-iodide symporter) (Na(+)/I(-) symporter) (Solute carrier family 5 member 5) Synthesized peptide derived from part region of human protein SC5A5 Monoclonal Antibody detects endogenous levels of protein. Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Reactivity Applications Gene Name Protein Name Immunogen Specificity Formulation	Human;Rat;Mouse; WB SLC5A5 NIS Sodium/iodide cotransporter (Na(+)/I(-) cotransporter) (Sodium-iodide symporter) (Na(+)/I(-) symporter) (Solute carrier family 5 member 5) Synthesized peptide derived from part region of human protein SC5A5 Monoclonal Antibody detects endogenous levels of protein.
Applications Gene Name Protein Name Immunogen Specificity Formulation	WB SLC5A5 NIS Sodium/iodide cotransporter (Na(+)/I(-) cotransporter) (Sodium-iodide symporter) (Na(+)/I(-) symporter) (Solute carrier family 5 member 5) Synthesized peptide derived from part region of human protein SC5A5 Monoclonal Antibody detects endogenous levels of protein.
Gene Name Protein Name Immunogen Specificity Formulation	SLC5A5 NIS Sodium/iodide cotransporter (Na(+)/I(-) cotransporter) (Sodium-iodide symporter) (Na(+)/I(-) symporter) (Solute carrier family 5 member 5) Synthesized peptide derived from part region of human protein SC5A5 Monoclonal Antibody detects endogenous levels of protein.
Protein Name Immunogen Specificity Formulation	Sodium/iodide cotransporter (Na(+)/I(-) cotransporter) (Sodium-iodide symporter) (Na(+)/I(-) symporter) (Solute carrier family 5 member 5) Synthesized peptide derived from part region of human protein SC5A5 Monoclonal Antibody detects endogenous levels of protein.
Immunogen Specificity Formulation	Synthesized peptide derived from part region of human protein SC5A5 Monoclonal Antibody detects endogenous levels of protein.
Specificity Formulation	SC5A5 Monoclonal Antibody detects endogenous levels of protein.
Formulation	· · · · · · · · · · · · · · · · · · ·
	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	
	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	70kD
Cell Pathway	Membrane; Multi-pass membrane protein.
Tissue Specificity	Expression is primarily in thyroid tissue, but also to a lower extent in mammary gland and ovary. Expression is reduced in tumors.
Function	disease:Defects in SLC5A5 are the cause of congenital hypothyroidism due to dyshormonogenesis type 1 (CHDH1) [MIM:274400]. CHDH1 is characterized by an inability of the thyroid to maintain a concentration difference of readily exchangeable iodine between the plasma and the thyroid gland, leading to congenital hypothyroidism.,function:Mediates iodide uptake in the thyroid
	gland.,similarity:Bélongs to the sodium:solute symporter (SSF) (TC 2.A.21) family.,tissue specificity:Expression is primarily in thyroid tissue, but also to a lower extent in mammary gland and ovary. Expression is reduced in tumors.,



UpingBio technology Co.,Ltd





matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images