



# SYT3 Monoclonal Antibody

Catalog No	YP-mAb-06258
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB
Gene Name	SYT3
Protein Name	Synaptotagmin-3 (Synaptotagmin III) (SytlIII)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	SYT3 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	64kD
Cell Pathway	Cell membrane ; Single-pass membrane protein . Cytoplasmic vesicle, secretory vesicle membrane ; Single-pass membrane protein .
Tissue Specificity	Expressed in melanocytes (PubMed:23999003).
Function	cofactor: Binds 3 calcium ions per subunit. The ions are bound to the C2 domains.,domain: The first C2 domain mediates Ca(2+)-dependent phospholipid binding.,function: May be involved in Ca(2+)-dependent exocytosis of secretory vesicles through Ca(2+) and phospholipid binding to the C2 domain or may serve as Ca(2+) sensors in the process of vesicular trafficking and exocytosis.,similarity: Belongs to the synaptotagmin family.,similarity: Contains 2 C2 domains.,subunit: Homodimer. Can also form heterodimers.,
Background	cofactor: Binds 3 calcium ions per subunit. The ions are bound to the C2 domains.,domain: The first C2 domain mediates Ca(2+)-dependent phospholipid binding.,function: May be involved in Ca(2+)-dependent exocytosis of secretory vesicles through Ca(2+) and phospholipid binding to the C2 domain or may serve as Ca(2+) sensors in the process of vesicular trafficking and exocytosis.,similarity: Belongs to the synaptotagmin family.,similarity: Contains 2 C2 domains.,subunit: Homodimer. Can also form heterodimers.,

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**