



TBD2A Monoclonal Antibody

Catalog No	YP-mAb-06268
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	TBC1D2 PARIS1 PP8997 TBC1D2A
Protein Name	TBC1 domain family member 2A (Armus) (Prostate antigen recognized and identified by SEREX 1) (PARIS-1)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	TBD2A Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	102kD
Cell Pathway	Cytoplasm . Cytoplasmic vesicle . Cell junction .
Tissue Specificity	Expressed in a broad range of tissues, especially in kidney, liver, lung and placenta. Also expressed in keratinocytes and epithelia-containing organs. Isoform 2 is differentially expressed in prostate normal and cancer cells (at protein level).
Function	function:May act as a GTPase-activating protein for Rab family protein(s).,similarity:Contains 1 PH domain.,similarity:Contains 1 Rab-GAP TBC domain.,tissue specificity:Expressed in a broad range of tissues. Differentially expressed in prostate normal and cancer cells.,
Background	function:May act as a GTPase-activating protein for Rab family protein(s).,similarity:Contains 1 PH domain.,similarity:Contains 1 Rab-GAP TBC domain.,tissue specificity:Expressed in a broad range of tissues. Differentially expressed in prostate normal and cancer cells.,
matters needing attention	Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images