



CENPT Monoclonal Antibody

Catalog No	YP-mAb-06453
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	CENPT C16orf56 ICEN22
Protein Name	Centromere protein T (CENP-T) (Interphase centromere complex protein 22)
Immunogen	Synthesized peptide derived from human protein . at AA range: 20-100
Specificity	CENPT Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	61kD
Cell Pathway	Nucleus. Chromosome, centromere . Chromosome, centromere, kinetochore . Constitutively localizes to centromeres throughout the cell cycle, and to kinetochores during mitosis. Localizes to the inner kinetochore, and may connect it to the outer kinetochore via its N-terminus. .
Tissue Specificity	Brain,Epithelium,Lung,Uterus,
Function	function:Component of the CENPA-NAC (nucleosome-associated) complex, a complex that plays a central role in assembly of kinetochore proteins, mitotic progression and chromosome segregation. The CENPA-NAC complex recruits the CENPA-CAD (nucleosome distal) complex and may be involved in incorporation of newly synthesized CENPA into centromeres.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,subcellular location:Localizes exclusively in the kinetochore domain of centromeres.,subunit:Component of the CENPA-NAC complex, at least composed of CENPA, CENPC, CENPH, CENPM, CENPN, CENPT and MLF1IP/CENPU. The CENPA-NAC complex interacts with the CENPA-CAD complex, composed of CENPI, CENPK, CENPL, CENPO, CENPP, CENPQ, CENPR and CENPS.,

**Background**

The centromere is a specialized chromatin domain, present throughout the cell cycle, that acts as a platform on which the transient assembly of the kinetochore occurs during mitosis. All active centromeres are characterized by the presence of long arrays of nucleosomes in which CENPA (MIM 117139) replaces histone H3 (see MIM 601128). CENPT is an additional factor required for centromere assembly (Foltz et al., 2006 [PubMed 16622419]).[supplied by OMIM, Mar 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images