



# M3K12 Monoclonal Antibody

Catalog No	YP-mAb-06462
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	MAP3K12 ZPK
Protein Name	Mitogen-activated protein kinase kinase kinase 12 (EC 2.7.11.25) (Dual leucine zipper bearing kinase) (DLK) (Leucine-zipper protein kinase) (ZPK) (MAPK-upstream kinase) (MUK) (Mixed lineage kinase)
Immunogen	Synthesized peptide derived from human protein . at AA range: 730-810
Specificity	M3K12 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	94kD
Cell Pathway	Cytoplasm . Cell membrane . Behaves essentially as an integral membrane protein. .
Tissue Specificity	Highly expressed in brain and kidney.
Function	catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,domain:Interacts with MBIP through the leucine-zipper motif.,function:May be an activator of the JNK/SAPK pathway. Phosphorylates beta-casein, histone 1 and myelin basic protein in vitro.,PTM:Autophosphorylated on Ser/Thr. Phosphorylated in cytosol under basal conditions and dephosphorylated when membrane-associated.,similarity:Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. MAP kinase kinase kinase subfamily.,similarity:Contains 1 protein kinase domain.,subunit:Interacts with MBIP.,tissue specificity:Highly expressed in brain and kidney.,
Background	This gene encodes a member of the serine/threonine protein kinase family. This kinase contains a leucine-zipper domain and is predominately expressed in neuronal cells. The phosphorylation state of this kinase in synaptic terminals was shown to be regulated by membrane depolarization via calcineurin. This kinase



forms heterodimers with leucine zipper containing transcription factors, such as cAMP responsive element binding protein (CREB) and MYC, and thus may play a regulatory role in PKA or retinoic acid induced neuronal differentiation. Alternatively spliced transcript variants encoding different proteins have been described.[provided by RefSeq, Jul 2010],

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images