





SNPC4 Monoclonal Antibody

YP-mAb-06502
IgG
Human;Rat;Mouse;
WB
SNAPC4 SNAP190
snRNA-activating protein complex subunit 4 (SNAPc subunit 4) (Proximal sequence element-binding transcription factor subunit alpha) (PSE-binding factor subunit alpha) (PTF subunit alpha) (snRNA-activa
Synthesized peptide derived from part region of human protein
SNPC4 Monoclonal Antibody detects endogenous levels of protein.
Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Monoclonal, Mouse,IgG
The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
WB 1:500-1:2000
1 mg/ml
≥90%
-20°C/1 year
161kD
Nucleus .
Epithelium,Teratocarcinoma,
function:Part of the SNAPc complex required for the transcription of both RNA polymerase II and III small-nuclear RNA genes. Binds to the proximal sequence element (PSE), a non-TATA-box basal promoter element common to these 2 types of genes. Recruits TBP and BRF2 to the U6 snRNA TATA box., similarity:Contains 2 Myb-like domains., similarity:Contains 3 HTH myb-type DNA-binding domains., subunit:Part of the SNAPc complex composed of 5 subunits: SNAPC1, SNAPC2, SNAPC3, SNAPC4 and SNAPC5. SNAPC4 interacts with SNAPC1, SNAPC2, SNAPC5, BRF2 and TBP.,
This gene encodes the largest subunit of the small nuclear RNA-activating protein (SNAP) complex. The encoded protein contains a Myb DNA-binding domain, and is essential for RNA polymerase II and III polymerase transcription from small nuclear RNA promoters. A mutation in this gene is associated with



UpingBio technology Co.,Ltd





matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images