



# CYFP1 Monoclonal Antibody

<b>Catalog No</b>	YP-mAb-06514
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB
<b>Gene Name</b>	CYFIP1 KIAA0068
<b>Protein Name</b>	Cytoplasmic FMR1-interacting protein 1 (Specifically Rac1-associated protein 1) (Sra-1) (p140sra-1)
<b>Immunogen</b>	Synthesized peptide derived from human protein . at AA range: 980-1060
<b>Specificity</b>	CYFP1 Monoclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-1:2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	137kD
<b>Cell Pathway</b>	Cytoplasm . Cytoplasm, perinuclear region . Cell projection, lamellipodium . Cell projection, ruffle . Cell junction, synapse, synaptosome . Highly expressed in the perinuclear region (By similarity). Enriched in synaptosomes (By similarity). Also enriched in membrane ruffles and at the tips of lamellipodia (By similarity). .
<b>Tissue Specificity</b>	B-cell lymphoma,Bone marrow,Brain,Hepatoma,Lung,Placenta,
<b>Function</b>	function:Involved in formation of membrane ruffles and lamellipodia protrusions and in axon outgrowth. Binds to F-actin but not to RNA.,similarity:Belongs to the CYFIP family.,subcellular location:Highly expressed in the perinuclear region. Enriched in synaptosomes, membrane ruffles and at the tips of lamellipodia.,subunit:Interacts with the active GTP-bound form of RAC1. Interacts with FMR1/FMRP but does not bind to related proteins FXR1 or FXR2. Component of the WAVE2 complex composed of ABI1, CYFIP1/SRA1, NCKAP1/NAP1 and WASF2/WAVE2 (By similarity). Interacts through its C-terminus with the C-terminus of DPYSL2/CRMP2 which is necessary for DPYSL2-induced axon outgrowth.,
<b>Background</b>	function:Involved in formation of membrane ruffles and lamellipodia protrusions and in axon outgrowth. Binds to F-actin but not to RNA.,similarity:Belongs to the CYFIP family.,subcellular location:Highly expressed in the perinuclear region.



Enriched in synaptosomes, membrane ruffles and at the tips of lamellipodia.,subunit:Interacts with the active GTP-bound form of RAC1. Interacts with FMR1/FMRP but does not bind to related proteins FXR1 or FXR2. Component of the WAVE2 complex composed of ABI1, CYFIP1/SRA1, NCKAP1/NAP1 and WASF2/WAVE2 (By similarity). Interacts through its C-terminus with the C-terminus of DPYSL2/CRMP2 which is necessary for DPYSL2-induced axon outgrowth.,

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images