



KDM5B Monoclonal Antibody

Catalog No	YP-mAb-06640
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	KDM5B JARID1B PLU1 RBBP2H1
Protein Name	Lysine-specific demethylase 5B (EC 1.14.11.-) (Cancer/testis antigen 31) (CT31) (Histone demethylase JARID1B) (Jumonji/ARID domain-containing protein 1B) (PLU-1) (Retinoblastoma-binding protein 2 homo
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	KDM5B Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	169kD
Cell Pathway	Nucleus .
Tissue Specificity	Ubiquitously expressed, with highest levels in testis. Down-regulated in melanoma and glioblastoma. Up-regulated in breast cancer (at protein level).
Function	cofactor:Fe(2+).,domain:Both the JmjC domain and the JmjN domain are required for enzymatic activity.,domain:The 2 first PHD-type zinc finger domains are required for transcription repression activity.,function:Histone demethylase that demethylates 'Lys-4' of histone H3, thereby playing a central role in histone code. Does not demethylate histone H3 'Lys-9' or H3 'Lys-27'. Demethylates trimethylated, dimethylated and monomethylated H3 'Lys-4'. Acts as a transcriptional corepressor for FOXG1B and PAX9. Favors the proliferation of breast cancer cells by repressing tumor suppressor genes such as BRCA1 and HOXA5. In contrast, may act as a tumor suppressor for melanoma.,similarity:Belongs to the JARID1 histone demethylase family.,similarity:Contains 1 ARID domain.,similarity:Contains 1 JmjC domain.,similarity:Contains 1 JmjN domain.,similarity:Contains 3 PHD-type zinc fingers.,subunit:Interac

**Background**

This gene encodes a lysine-specific histone demethylase that belongs to the jumonji/ARID domain-containing family of histone demethylases. The encoded protein is caMABLE of demethylating tri-, di- and monomethylated lysine 4 of histone H3. This protein plays a role in the transcriptional repression or certain tumor suppressor genes and is upregulated in certain cancer cells. This protein may also play a role in genome stability and DNA repair. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Sep 2015],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images