





## **TBD Monoclonal Antibody**

| Catalog No         | YP-mAb-06660   |
|--------------------|--|
| Isotype            | IgG  |
| Reactivity         | Human;Rat;Mouse;   |
| Applications       | WB   |
| Gene Name          | TUBD1 TUBD   |
| Protein Name       | Tubulin delta chain (Delta-tubulin)  |
| Immunogen          | Synthesized peptide derived from part region of human protein  |
| Specificity        | TBD Monoclonal Antibody detects endogenous levels of protein.  |
| Formulation        | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.   |
| Source             | Monoclonal, Mouse,IgG  |
| Purification       | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   |
| Dilution           | WB 1:500-1:2000  |
| Concentration      | 1 mg/ml  |
| Purity             | ≥90%   |
| Storage Stability  | -20°C/1 year   |
| Synonyms           |  |
| Observed Band      | 49kD   |
| Cell Pathway       | Nucleus . Cytoplasm . Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole . Cell projection, cilium . Associated with centrioles. Both cytoplasmic and nuclear. In the elongating spermatid it is associated with the manchette, a specialized microtubule system present during reshaping of the sperm head.                          |
| Tissue Specificity | Embryo,Eye,  |
| Function           | alternative products:Experimental confirmation may be lacking for some isoforms,function:In the elongating spermatid it is associated with the manchette, a specialized microtubule system present during reshaping of the sperm head.,similarity:Belongs to the tubulin family.,subcellular location:Associated with centrioles. Both cytoplasmic and nuclear., |
| Background         | alternative products:Experimental confirmation may be lacking for some isoforms,function:In the elongating spermatid it is associated with the manchette, a specialized microtubule system present during reshaping of the sperm head.,similarity:Belongs to the tubulin family.,subcellular location:Associated with centrioles. Both cytoplasmic and nuclear., |



## UpingBio technology Co.,Ltd





matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**