





CLK3 Monoclonal Antibody

YP-mAb-06731 IgG
•
Human;Mouse;Rat
WB
CLK3
Dual specificity protein kinase CLK3 (EC 2.7.12.1) (CDC-like kinase 3)
Synthesized peptide derived from part region of human protein
CLK3 Monoclonal Antibody detects endogenous levels of protein.
Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Monoclonal, Mouse,IgG
The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
WB 1:500-1:2000
1 mg/ml
≥90%
-20°C/1 year
70kD
[Isoform 1]: Nucleus. Cytoplasm . Cytoplasmic vesicle, secretory vesicle, acrosome .; [Isoform 2]: Nucleus speckle. Co-localizes with serine- and arginine-rich (SR) proteins in the nuclear speckles.
Endothelial cells.
catalytic activity:ATP + a protein = ADP + a phosphoprotein.,function:Phosphorylates serine- and arginine-rich (SR) proteins of the spliceosomal complex. May be a constituent of a network of regulatory mechanisms that enable SR proteins to control RNA splicing. Phosphorylates serines, threonines and tyrosines.,PTM:Autophosphorylates on all three types of residues.,similarity:Belongs to the protein kinase superfamily.,similarity:Belongs to the protein kinase family. Lammer subfamily.,similarity:Contains 1 protein kinase domain.,
CDC like kinase 3(CLK3) Homo sapiens This gene encodes a protein belonging to the serine/threonine type protein kinase family. This protein is a nuclear dual-specificity kinase that regulates the intranuclear distribution of the serine/arginine-rich (SR) family of splicing factors. Two transcript variants encoding different isoforms have been found for this gene. Related pseudogenes



UpingBio technology Co.,Ltd





matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images