





## NEK3 Monoclonal Antibody

Catalog No         YP-mAb-06737           Isotype         IgG           Reactivity         Mouse           Applications         WB           Gene Name         NEK3           Protein Name         Serine/fthreonine-protein kinase Nek3 (EC 2.7.11.1) (HSPK 36) (Never in mitosis A-related kinase 3) (NimA-related protein kinase 3)           Immunogen         Synthesized peptide derived from part region of human protein           Specificity         NEK3 Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-1:2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         55kD           Cell Pathway         Cytoplasm . Cell projection, axon .           Tissue Specificity         Up-regulated in malignant versus normal breast tissue. Isoform 2 shows a high level of expression in testis, ovary and brain.           Function         catalytic activity:ATP + a protein = ADP + a phosphoprotein, catacor Magnesium, function:Kinase that ma		
Reactivity         Mouse           Applications         WB           Gene Name         NEK3           Protein Name         Serine/threonine-protein kinase Nek3 (EC 2.7.11.1) (HSPK 36) (Never in mitosis A-related kinase 3) (NimA-related protein kinase 3)           Immunogen         Synthesized peptide derived from part region of human protein           Specificity         NEK3 Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-1:2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         55kD           Cell Pathway         Cytoplasm . Cell projection, axon .           Tissue Specificity         Up-regulated in malignant versus normal breast tissue. Isoform 2 shows a high level of expression in testis, ovary and brain.           Function         catalytic activity.ATP + a protein = ADP + a protein kinase more protein, cofactor, Magnesium, function, Kinase that may play a role in mitotic regulation, similarity. Edongs to the protein kinase superfamily, NEK Ser/Thr protein kinase family.	Catalog No	YP-mAb-06737
Applications WB  Gene Name NEK3  Protein Name Serine/threonine-protein kinase Nek3 (EC 2.7.11.1) (HSPK 36) (Never in mitosis A-related kinase 3) (NimA-related protein kinase 3)  Immunogen Synthesized peptide derived from part region of human protein  Specificity NEK3 Monoclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 55kD  Cell Pathway Cytoplasm . Cell projection, axon .  Tissue Specificity Up-regulated in malignant versus normal breast tissue. Isoform 2 shows a high level of expression in testis, ovary and brein.  Function catalytic activity. ATP + a protein = ADP + a phosphoprotein, cofactor. Magnesium, function: Kinase that may play a role in mitotic regulation, similarity. Belongs to the protein kinase superfamily, similarity. Belongs to the protein kinase superfamily, similarity. Belongs to the protein kinase superfamily, nembers in that it is not cell cycle regulated and is found primarily in the cytoplasm. The kinase is activated by prolactin stimulation, leading to phosphorylation of VAV2 guanine nucleotide exchange factor, paxillin, and activation of the RAC1 GT Pase. Multiple transcript variants encoding different	Isotype	IgG
Gene Name         NEK3           Protein Name         Serine/threonine-protein kinase Nek3 (EC 2.7.11.1) (HSPK 36) (Never in mitosis A-related kinase 3) (NimA-related protein kinase 3)           Immunogen         Synthesized peptide derived from part region of human protein           Specificity         NEK3 Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-1:2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         55kD           Cell Pathway         Cytoplasm . Cell projection, axon .           Tissue Specificity         Up-regulated in malignant versus normal breast tissue. Isoform 2 shows a high level of expression in testis, ovary and brain.           Function         catalytic activity. ATP + a protein = ADP + a phosphoprotein . cofactor Magnesium , function: kinase that may play a role in mitotic requiation . similarity. Belongs to the protein kinase superfamily, NEK Ser/Thr protein kinase family . Nimilarity. Belongs to the protein kinase superfamily, similarity. The kinase is activated by prolactin stimulation, leading to phosphorylation of	Reactivity	Mouse
Protein Name   Serine/threonine-protein kinase Nek3 (EC 2.7.11.1) (HSPK 36) (Never in mitosis A-related kinase 3) (NimA-related protein kinase 3)	Applications	WB
Immunogen         Synthesized peptide derived from part region of human protein           Specificity         NEK3 Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-1:2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         55kD           Cell Pathway         Cytoplasm . Cell projection, axon .           Tissue Specificity         Up-regulated in malignant versus normal breast tissue. Isoform 2 shows a high level of expression in testis, ovary and brain.           Function         catalytic activity.ATP + a protein = ADP + a phosphoprotein. cofactor/Magnesium. function/kinase that may play a role in mitotic regulation, similarity. Belongs to the protein kinase superfamily. NEK Ser/Thr protein kinase superfamily. NEK Ser/Thr protein kinase family. NIMA subfamily, similarity Contains 1 protein kinase domain.           Background         This gene encodes a member of the NimA (never in mitosis A) family of serine/threonine protein kinases. The encoded protein differs from other NimA family members in that it is not cell cycle regulated and is found primarily in the cytoplasm. The kinase	Gene Name	NEK3
Specificity         NEK3 Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-1:2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band           Cell Pathway         Cytoplasm . Cell projection, axon .           Tissue Specificity         Up-regulated in malignant versus normal breast tissue. Isoform 2 shows a high level of expression in testis, ovary and brain.           Function         catalytic activity.ATP + a protein = ADP + a phosphoprotein, cofactor.Magnesium. function: kinase that may play a role in mitotic regulation, similarity. Belongs to the protein kinase superfamily. NEK Ser/Thr protein kinase family. NilMA subfamily., similarity. Contains 1 protein kinase domain.           Background         This gene encodes a member of the NimA (never in mitosis A) family of serine/threonine protein kinases. The encoded protein differs from other NimA family members in that it is not cell cycle regulated and is found primarily in the cytoplasm. The kinase is activated by prolacitin stimulation, leading to phosphorylation of the RACH GTPase. Multiple transcript variants encoding different	Protein Name	Serine/threonine-protein kinase Nek3 (EC 2.7.11.1) (HSPK 36) (Never in mitosis A-related kinase 3) (NimA-related protein kinase 3)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 55kD  Cell Pathway Cytoplasm · Cell projection, axon ·  Tissue Specificity Up-regulated in malignant versus normal breast tissue. Isoform 2 shows a high level of expression in testis, ovary and brain.  Function Catalytic activity:ATP + a protein = ADP + a phosphorptein, cofactor: Magnesium., function:Kinase that may play a role in mitotic regulation, similarity: Belongs to the protein kinase superfamily. NEK Ser/Thr protein kinase family. NIMA subfamily., similarity:Contains 1 protein kinase domain.,  This gene encodes a member of the NimA (never in mitosis A) family of serine/threonine protein kinases. The encoded protein differs from other NimA family members in that it is not cell cycle regulated and is found primarily in the cytoplasm. The kinase is activated by prolactin stimulation, leading to phosphorylation of VA/2 guanine nucleotide exchange factor, paxillin, and activation of the RAC1 GTPase. Multiple transcript variants encoding different	Immunogen	Synthesized peptide derived from part region of human protein
Source       Monoclonal, Mouse,IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-1:2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms         Observed Band       55kD         Cell Pathway       Cytoplasm . Cell projection, axon .         Tissue Specificity       Up-regulated in malignant versus normal breast tissue. Isoform 2 shows a high level of expression in testis, ovary and brain.         Function       catalytic activity:ATP + a protein = ADP + a phosphoprotein., cofactor:Magnesium., function:Kinase that may play a role in mitotic regulation., similarity:Belongs to the protein kinase superfamily. NEK Ser/Thr protein kinase superfamily. NIMA subfamily., similarity:Contains 1 protein kinase domain.         Background       This gene encodes a member of the NimA (never in mitosis A) family of serine/threonine protein kinases. The encoded protein differs from other NimA family members in that it is not cell cycle regulated and is found primarily in the cytoplasm. The kinase is activated by prolactin stimulation, leading to phosphorylation of VAV2 guanine nucleotide exchange factor, paxilin, and activation of the RAC1 GTPase. Multiple transcript variants encoding different	Specificity	NEK3 Monoclonal Antibody detects endogenous levels of protein.
Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-1:2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  Observed Band  55kD  Cell Pathway  Cytoplasm . Cell projection, axon .  Tissue Specificity  Up-regulated in malignant versus normal breast tissue. Isoform 2 shows a high level of expression in testis, ovary and brain.  Function  catalytic activity:ATP + a protein = ADP + a phosphoprotein, cofactor:Magnesium, function:Kinase that may play a role in mitotic regulation, similarity:Belongs to the protein kinase superfamily. NEK Ser/Thr protein kinase family. NIMA subfamily.,similarity:Contains 1 protein kinase domain.  Background  This gene encodes a member of the NimA (never in mitosis A) family of serine/threonine protein kinases. The encoded protein differs from other NimA family members in that it is not cell cycle regulated and is found primarily in the cytoplasm. The kinase is activated by prolactin stimulation, leading to phosphorylation of VAV2 guanine nucleotide exchange factor, paxillin, and activation of the RAC1 GTPase. Multiple transcript variants encoding different	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 55kD  Cell Pathway Cytoplasm . Cell projection, axon .  Tissue Specificity Up-regulated in malignant versus normal breast tissue. Isoform 2 shows a high level of expression in testis, ovary and brain.  Function catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,function:Kinase that may play a role in mitotic regulation.,similarity:Belongs to the protein kinase superfamily. NEK Ser/Thr protein kinase family. NIMA subfamily.,similarity:Contains 1 protein kinase domain.  Background This gene encodes a member of the NimA (never in mitosis A) family of serine/threonine protein kinases. The encoded protein differs from other NimA family members in that it is not cell cycle regulated and is found primarily in the cytoplasm. The kinase is activated by prolactin stimulation, leading to phosphorylation of VAV2 guanine nucleotide exchange factor, paxillin, and activation of the RAC1 GTPase. Multiple transcript variants encoding different	Source	Monoclonal, Mouse,lgG
Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band         Cell Pathway       Cytoplasm . Cell projection, axon .         Tissue Specificity       Up-regulated in malignant versus normal breast tissue. Isoform 2 shows a high level of expression in testis, ovary and brain.         Function       catalytic activity.ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,function:Kinase that may play a role in mitotic regulation.,similarity:Belongs to the protein kinase superfamily. NEK Ser/Thr protein kinase family. NIMA subfamily.,similarity:Contains 1 protein kinase domain.,         Background       This gene encodes a member of the NimA (never in mitosis A) family of serine/threonine protein kinases. The encoded protein differs from other NimA family members in that it is not cell cycle regulated and is found primarily in the cytoplasm. The kinase is activated by prolactin stimulation, leading to phosphorylation of VAV2 guanine nucleotide exchange factor, paxillin, and activation of the RAC1 GTPase. Multiple transcript variants encoding different	Purification	·
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 55kD  Cell Pathway Cytoplasm . Cell projection, axon .  Tissue Specificity Up-regulated in malignant versus normal breast tissue. Isoform 2 shows a high level of expression in testis, ovary and brain.  Function catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,function:Kinase that may play a role in mitotic regulation.,similarity:Belongs to the protein kinase superfamily. NEK Ser/Thr protein kinase family. NIMA subfamily.,similarity:Contains 1 protein kinase domain.,  Background This gene encodes a member of the NimA (never in mitosis A) family of serine/threonine protein kinases. The encoded protein differs from other NimA family members in that it is not cell cycle regulated and is found primarily in the cytoplasm. The kinase is activated by prolactin stimulation, leading to phosphorylation of VAV2 guanine nucleotide exchange factor, paxillin, and activation of the RAC1 GTPase. Multiple transcript variants encoding different	Dilution	WB 1:500-1:2000
Synonyms  Observed Band  Cell Pathway  Cytoplasm . Cell projection, axon .  Up-regulated in malignant versus normal breast tissue. Isoform 2 shows a high level of expression in testis, ovary and brain.  Function  Catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,function:Kinase that may play a role in mitotic regulation.,similarity:Belongs to the protein kinase superfamily. NEK Ser/Thr protein kinase family. NIMA subfamily.,similarity:Contains 1 protein kinase domain.,  Background  This gene encodes a member of the NimA (never in mitosis A) family of serine/threonine protein kinases. The encoded protein differs from other NimA family members in that it is not cell cycle regulated and is found primarily in the cytoplasm. The kinase is activated by prolactin stimulation, leading to phosphorylation of VAV2 guanine nucleotide exchange factor, paxillin, and activation of the RAC1 GTPase. Multiple transcript variants encoding different	Concentration	1 mg/ml
Synonyms  Observed Band 55kD  Cell Pathway Cytoplasm . Cell projection, axon .  Tissue Specificity Up-regulated in malignant versus normal breast tissue. Isoform 2 shows a high level of expression in testis, ovary and brain.  Function catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,function:Kinase that may play a role in mitotic regulation.,similarity:Belongs to the protein kinase superfamily. NEK Ser/Thr protein kinase family. NIMA subfamily.,similarity:Contains 1 protein kinase domain.,  Background This gene encodes a member of the NimA (never in mitosis A) family of serine/threonine protein kinases. The encoded protein differs from other NimA family members in that it is not cell cycle regulated and is found primarily in the cytoplasm. The kinase is activated by prolactin stimulation, leading to phosphorylation of VAV2 guanine nucleotide exchange factor, paxillin, and activation of the RAC1 GTPase. Multiple transcript variants encoding different	Purity	≥90%
Observed Band         55kD           Cell Pathway         Cytoplasm . Cell projection, axon .           Tissue Specificity         Up-regulated in malignant versus normal breast tissue. Isoform 2 shows a high level of expression in testis, ovary and brain.           Function         catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,function:Kinase that may play a role in mitotic regulation.,similarity:Belongs to the protein kinase superfamily. NEK Ser/Thr protein kinase family., similarity:Belongs to the protein kinase superfamily. NEK Ser/Thr protein kinase family. NIMA subfamily.,similarity:Contains 1 protein kinase domain.,           Background         This gene encodes a member of the NimA (never in mitosis A) family of serine/threonine protein kinases. The encoded protein differs from other NimA family members in that it is not cell cycle regulated and is found primarily in the cytoplasm. The kinase is activated by prolactin stimulation, leading to phosphorylation of VAV2 guanine nucleotide exchange factor, paxillin, and activation of the RAC1 GTPase. Multiple transcript variants encoding different	Storage Stability	-20°C/1 year
Cell Pathway  Cytoplasm . Cell projection, axon .  Up-regulated in malignant versus normal breast tissue. Isoform 2 shows a high level of expression in testis, ovary and brain.  Function  catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,function:Kinase that may play a role in mitotic regulation.,similarity:Belongs to the protein kinase superfamily. NEK Ser/Thr protein kinase family. NIMA subfamily.,similarity:Contains 1 protein kinase domain.,  Background  This gene encodes a member of the NimA (never in mitosis A) family of serine/threonine protein kinases. The encoded protein differs from other NimA family members in that it is not cell cycle regulated and is found primarily in the cytoplasm. The kinase is activated by prolactin stimulation, leading to phosphorylation of VAV2 guanine nucleotide exchange factor, paxillin, and activation of the RAC1 GTPase. Multiple transcript variants encoding different	Synonyms	
Tissue Specificity  Up-regulated in malignant versus normal breast tissue. Isoform 2 shows a high level of expression in testis, ovary and brain.  Function  catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,function:Kinase that may play a role in mitotic regulation.,similarity:Belongs to the protein kinase superfamily. NEK Ser/Thr protein kinase family. NIMA subfamily.,similarity:Contains 1 protein kinase domain.,  This gene encodes a member of the NimA (never in mitosis A) family of serine/threonine protein kinases. The encoded protein differs from other NimA family members in that it is not cell cycle regulated and is found primarily in the cytoplasm. The kinase is activated by prolactin stimulation, leading to phosphorylation of VAV2 guanine nucleotide exchange factor, paxillin, and activation of the RAC1 GTPase. Multiple transcript variants encoding different	Observed Band	55kD
Level of expression in testis, ovary and brain.	Cell Pathway	Cytoplasm . Cell projection, axon .
phosphoprotein.,cofactor:Magnesium.,function:Kinase that may play a role in mitotic regulation.,similarity:Belongs to the protein kinase superfamily. NEK Ser/Thr protein kinase family. NIMA subfamily.,similarity:Contains 1 protein kinase domain.,  Background  This gene encodes a member of the NimA (never in mitosis A) family of serine/threonine protein kinases. The encoded protein differs from other NimA family members in that it is not cell cycle regulated and is found primarily in the cytoplasm. The kinase is activated by prolactin stimulation, leading to phosphorylation of VAV2 guanine nucleotide exchange factor, paxillin, and activation of the RAC1 GTPase. Multiple transcript variants encoding different	Tissue Specificity	
serine/threonine protein kinases. The encoded protein differs from other NimA family members in that it is not cell cycle regulated and is found primarily in the cytoplasm. The kinase is activated by prolactin stimulation, leading to phosphorylation of VAV2 guanine nucleotide exchange factor, paxillin, and activation of the RAC1 GTPase. Multiple transcript variants encoding different	Function	phosphoprotein.,cofactor:Magnesium.,function:Kinase that may play a role in mitotic regulation.,similarity:Belongs to the protein kinase superfamily. NEK Ser/Thr protein kinase family. NIMA subfamily.,similarity:Contains 1 protein kinase
	Background	serine/threonine protein kinases. The encoded protein differs from other NimA family members in that it is not cell cycle regulated and is found primarily in the cytoplasm. The kinase is activated by prolactin stimulation, leading to phosphorylation of VAV2 guanine nucleotide exchange factor, paxillin, and activation of the RAC1 GTPase. Multiple transcript variants encoding different



## UpingBio technology Co.,Ltd





matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**