

KGP1 Monoclonal Antibody

Catalog No	YP-mAb-06745
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	PRKG1 PRKG1B PRKGR1A PRKGR1B
Protein Name	cGMP-dependent protein kinase 1 (cGK 1) (cGK1) (EC 2.7.11.12) (cGMP-dependent protein kinase I) (cGKI)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	KGP1 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	73kD
Cell Pathway	Cytoplasm . Colocalized with TRPC7 in the plasma membrane
Tissue Specificity	Primarily expressed in lung and placenta.
Function	catalytic activity:ATP + a protein = ADP + a phosphoprotein.,similarity:Belongs to the protein kinase superfamily.,
Background	Mammals have three different isoforms of cyclic GMP-dependent protein kinase (lalpha, lbeta, and II). These PRKG isoforms act as key mediators of the nitric oxide/cGMP signaling pathway and are important components of many signal transduction processes in diverse cell types. This PRKG1 gene on human chromosome 10 encodes the soluble lalpha and lbeta isoforms of PRKG by alternative transcript splicing. A separate gene on human chromosome 4, PRKG2 encodes the membrane-bound PRKG isoform II. The PRKG1 proteins play a central role in regulating cardiovascular and neuronal functions in addition to relaxing smooth muscle tone, preventing platelet aggregation, and modulating ce growth. This gene is most strongly expressed in all types of smooth muscle, platelets, cerebellar Purkinje cells, hippocampal neurons, and the lateral amygdala. Isoforms lalpha and Ibeta have identical cGMP-bindin



UpingBio technology Co.,Ltd





matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images