

SPRL1 Monoclonal Antibody

lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in		
Reactivity Human;Rat;Mouse Applications WB Gene Name SPARCL1 Protein Name SPARC-like protein 1 (High endothelial venule protein) (Hevin) (MAST 9) Immunogen Synthesized peptide derived from part region of human protein Specificity SPRL1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 73kD Cell Pathway Secreted, extracellular space, extracellular matrix. Tissue Specificity Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes. Function similarity:Belongs to the SPARC family, similarity:Contains 1 EF-hand domain, similarity:Contains 1 follistatin-like domain, similarity:Contains 1 large and peripheral blood leukocytes. Background similarity skelengs to SPARC family, similarity:Contains 1 EF-hand domain, similarity:Contains 1 splacenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood domain, similarity:Contains 1 EF-hand domain, similarity:Contains 1 splacenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes. Background similarity:Contains 1 follistatin-like domain, similarity:Contains 1 EF-hand domain, similarity:Contains 1 follistatin-like domain, and no expression in kidney, liver, and peripheral blood leukocytes.	Catalog No	YP-mAb-06912
Applications WB Gene Name SPARCL1 Protein Name SPARC-like protein 1 (High endothelial venule protein) (Hevin) (MAST 9) Immunogen Synthesized peptide derived from part region of human protein Specificity SPRL1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 73kD Cell Pathway Secreted, extracellular space, extracellular matrix. Tissue Specificity Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and domain, similarity-Contains 1 foliastain-like domain, similarity-Contains 1 Kazal-like domain, tissue specificity; Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes. Function similarity-Contains 1 foliastatin-like domain, similarity-Contains 1 (Kazal-like domain, tissue specificity; Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes. Background	Isotype	IgG
Gene Name SPARCLIke protein 1 (High endothelial venule protein) (Hevin) (MAST 9) Immunogen Synthesized peptide derived from part region of human protein Specificity SPRL1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dillution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 73kD Cell Pathway Secreted, extracellular space, extracellular matrix. Tissue Specificity Tissue Specificity Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes. Function similarity: Belongs to the SPARC family, similarity: Contains 1 follstatin-like domain, similarity: Contains	Reactivity	Human;Rat;Mouse
Protein Name SPARC-like protein 1 (High endothelial venule protein) (Hevin) (MAST 9) Immunogen Synthesized peptide derived from part region of human protein Specificity SPRL1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Secreted, extracellular space, extracellular matrix. Tissue Specificity Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes. Function similarity:Belongs to the SPARC family, similarity:Contains 1 EF-hand domain, similarity:Contains 1 follistatin-like domain, similarity:Contains 1 kazal-like domain, similarity:Contains 1 follistatin-like domain, si	Applications	WB
Immunogen Synthesized peptide derived from part region of human protein Specificity SPRL1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 73kD Cell Pathway Secreted, extracellular space, extracellular matrix. Tissue Specificity Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and trymus, and no expression in kidney, liver, and peripheral blood leukocytes. Function similarity:Belongs to the SPARC family, similarity:Contains 1 EF-hand domain, similarity:Contains 1 follistatin-like domain, similarity:Contains 1 Kazal-like domain, similarity:Contains 1 follistatin-like domain, si	Gene Name	SPARCL1
Specificity SPRL1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 73kD Cell Pathway Secreted, extracellular space, extracellular matrix. Tissue Specificity Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes. Function similarity:Belongs to the SPARC family, similarity:Contains 1 EF-hand domain., similarity:Contains 1 follistatin-like domain., similarity:Contains 1 kazal-like domain, simal intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes. Background similarity:Belongs to the SPARC family, similarity:Contains 1 EF-hand domain., similarity:Belongs to the SPARC family, similarity:Contains 1 EF-hand domain., similarity:Contains 1 follistatin-like domain, similarity:Contains 1 Kazal-like domain., similarity:Contains 1 follistatin-like domain., similarity:Contains 1 follistatin-like domain, similarity:Conta	Protein Name	SPARC-like protein 1 (High endothelial venule protein) (Hevin) (MAST 9)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 73kD Cell Pathway Secreted, extracellular space, extracellular matrix. Tissue Specificity Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes. Function similarity:Belongs to the SPARC familysimilarity:Contains 1 EF-hand domain.,similarity:Contains 1 follistatin-like domain.,similarity:Contains 1 kazal-like domain.,itssue specificity:Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, and creas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes., spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes., spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes., spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes., spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes.	Immunogen	Synthesized peptide derived from part region of human protein
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 73kD Cell Pathway Secreted, extracellular space, extracellular matrix. Tissue Specificity Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes. Function similarity:Belongs to the SPARC family, similarity:Contains 1 EF-hand domain, similarity:Contains 1 follistatin-like domain, similarity:Contains 1 kazal-like domain, sissule specificity:Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes. Background similarity:Belongs to the SPARC family, similarity:Contains 1 EF-hand domain. similarity:Contains 1 follistatin-like domain, similarity:Contains 1 kazal-like domain, sissule specificity-Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney,	Specificity	SPRL1 Monoclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 73kD Cell Pathway Secreted, extracellular space, extracellular matrix. Tissue Specificity Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes. Function Similarity:Belongs to the SPARC family, similarity:Contains 1 EF-hand domain., similarity:Contains 1 Kazal-like domain., tissue specificity:Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes. Function Similarity:Belongs to the SPARC family, similarity:Contains 1 EF-hand domain, similarity:Contains 1 (Section and peripheral blood leukocytes.) Background Similarity:Belongs to the SPARC family, similarity:Contains 1 EF-hand domain, similarity:Contains 1 (Kazal-like domain, similarity:Contains	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 73kD Cell Pathway Secreted, extracellular space, extracellular matrix. Tissue Specificity Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes. Function similarity: Belongs to the SPARC family, similarity: Contains 1 EF-hand domain., similarity: Contains 1 follistatin-like domain., similarity: Contains 1 kazal-like domain., tissue specificity: Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes. Background domain., similarity: Contains 1 follistatin-like domain., similarity: Contains 1 Kazal-like domain., sissue specificity: Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes.	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 73kD Cell Pathway Secreted, extracellular space, extracellular matrix. Tissue Specificity Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes. Function similarity:Belongs to the SPARC familysimilarity:Contains 1 EF-hand domain.,similarity:Contains 1 follistatin-like domain.,similarity:Contains 1 Kazal-like domain. tissue specificity:Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes. Background similarity:Belongs to the SPARC familysimilarity:Contains 1 EF-hand domain.,similarity:Contains 1 follistatin-like domain.,similarity:Contains 1 Kazal-like domain.,similarity:Contains 1 kazal-like domain.,similarity:Contains 1 follistatin-like domain.,similarity:Contains 1 kazal-like dom	Purification	·
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 73kD Cell Pathway Secreted, extracellular space, extracellular matrix. Tissue Specificity Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes. Function similarity:Belongs to the SPARC family.,similarity:Contains 1 EF-hand domain.,similarity:Contains 1 follistatin-like domain.,similarity:Contains 1 Kazal-like domain.,tissue specificity:Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes. Background similarity:Belongs to the SPARC family.,similarity:Contains 1 EF-hand domain.,similarity:Contains 1 follistatin-like domain.,similarity:Contains 1 Kazal-like domain.,tissue specificity:Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and locenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and locenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and locenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and locenta pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and literature.	Dilution	WB 1:500-1:2000
Synonyms Observed Band 73kD Cell Pathway Secreted, extracellular space, extracellular matrix. Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes. Function similarity:Belongs to the SPARC family.,similarity:Contains 1 EF-hand domain.,similarity:Contains 1 follistatin-like domain.,similarity:Contains 1 Kazal-like domain.,tissue specificity:Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes., similarity:Belongs to the SPARC family.,similarity:Contains 1 EF-hand domain.,similarity:Contains 1 follistatin-like domain.,similarity:Contains 1 Kazal-like domain.,tissue specificity:Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes.	Concentration	1 mg/ml
Synonyms Observed Band 73kD Cell Pathway Secreted, extracellular space, extracellular matrix. Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes. Function similarity:Belongs to the SPARC family.,similarity:Contains 1 EF-hand domain.,similarity:Contains 1 follistatin-like domain.,similarity:Contains 1 Kazal-like domain.,tissue specificity:Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes., Background similarity:Belongs to the SPARC family.,similarity:Contains 1 EF-hand domain.,similarity:Contains 1 follistatin-like domain.,similarity:Contains 1 Kazal-like domain.,tissue specificity:Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver,	Purity	≥90%
Cell Pathway Secreted, extracellular space, extracellular matrix. Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes. Function similarity:Belongs to the SPARC family.,similarity:Contains 1 EF-hand domain.,similarity:Contains 1 follistatin-like domain.,similarity:Contains 1 Kazal-like domain.,tissue specificity:Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes., Background similarity:Belongs to the SPARC family.,similarity:Contains 1 EF-hand domain.,similarity:Contains 1 follistatin-like domain.,similarity:Contains 1 Kazal-like domain.,tissue specificity:Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver,	Storage Stability	-20°C/1 year
Cell Pathway Secreted, extracellular space, extracellular matrix. Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes. Function similarity:Belongs to the SPARC family.,similarity:Contains 1 EF-hand domain.,similarity:Contains 1 follistatin-like domain.,similarity:Contains 1 Kazal-like domain.,tissue specificity:Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes., Similarity:Belongs to the SPARC family.,similarity:Contains 1 EF-hand domain.,similarity:Contains 1 flistatin-like domain.,similarity:Contains 1 Kazal-like domain.,tissue specificity:Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver,	Synonyms	
Tissue Specificity Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes. Function similarity:Belongs to the SPARC family.,similarity:Contains 1 EF-hand domain.,similarity:Contains 1 follistatin-like domain.,similarity:Contains 1 Kazal-like domain.,tissue specificity:Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes., Similarity:Belongs to the SPARC family.,similarity:Contains 1 EF-hand domain.,similarity:Contains 1 follistatin-like domain.,similarity:Contains 1 Kazal-like domain.,tissue specificity:Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver,	Observed Band	73kD
intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes. Function similarity:Belongs to the SPARC family.,similarity:Contains 1 EF-hand domain.,similarity:Contains 1 follistatin-like domain.,similarity:Contains 1 Kazal-like domain.,tissue specificity:Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes., similarity:Belongs to the SPARC family.,similarity:Contains 1 EF-hand domain.,similarity:Contains 1 follistatin-like domain.,similarity:Contains 1 Kazal-like domain.,tissue specificity:Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver,	Cell Pathway	Secreted, extracellular space, extracellular matrix .
domain.,similarity:Contains 1 follistatin-like domain.,similarity:Contains 1 Kazal-like domain.,tissue specificity:Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver, and peripheral blood leukocytes., Background similarity:Belongs to the SPARC family.,similarity:Contains 1 EF-hand domain.,similarity:Contains 1 follistatin-like domain.,similarity:Contains 1 Kazal-like domain.,tissue specificity:Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver,	Tissue Specificity	intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and
domain.,similarity:Contains 1 follistatin-like domain.,similarity:Contains 1 Kazal-like domain.,tissue specificity:Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver,	Function	domain.,similarity:Contains 1 follistatin-like domain.,similarity:Contains 1 Kazal-like domain.,tissue specificity:Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver,
	Background	domain.,similarity:Contains 1 follistatin-like domain.,similarity:Contains 1 Kazal-like domain.,tissue specificity:Highly expressed in lymph node, brain, heart, lung, skeletal muscle, ovary, small intestine, and colon, with lower levels in placenta, pancreas, testis, spleen, and thymus, and no expression in kidney, liver,



UpingBio technology Co.,Ltd





matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images