



PP2BB Monoclonal Antibody

IgG		
Applications WB Gene Name PPP3CB CALNA2 CALNB CNA2 Protein Name 3.1.3.16) (CAM—PRP catalytic subunit) (Calmodulin-dependent calcineurin A subunit beta isoform) Synthesized peptide derived from part region of human protein Specificity PP2BB Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 57kD Cell Pathway Cytoplasm Treatocarcinoma, Testis, alternative products: Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative spiloning events, catalytic activity. A phosphoprotein + H(2)C a protein + phosphate, cofactor:Binds 1 Fe(3+) ion per subunit, cofactor:Binds 1 Fe(3+) ion per subunit, colactor:Binds 2 This subunit may have a role in the calmodulin activation of calcineurin, similarity. Belongs to the PPP phosphatese family. PP-2B subfamily, subunit Composed of two components (A and B), the A component is the catalytic subunit and the B component confers calcium sensitivity. alternative sproducts: Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from prosphatase This subunit may have a role in the calmodulin activation of calcineurin, similarity. Belongs to the PPP phosphatese family. PP-2B subfamily, subunit Composed of two components (A and B), the A component is the catalytic subunit and the B component confers calcium sensitivity. alternative products: Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events catalytic activity. A phosphoprotein + H(2)C = a protein +	Catalog No	YP-mAb-06919
Applications WB Gene Name PPP3CB CALNA2 CALNB CNA2 Serine/threonine-protein phosphatase 2B catalytic subunit beta isoform (EC 3.1.3.16) (CAM-PRP catalytic subunit) (Calmodulin-dependent calcineurin A subunit beta isoform) Synthesized peptide derived from part region of human protein Specificity PP2BB Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml 290% Storage Stability -20°C/1 year Synonyms Observed Band 57kD Cell Pathway Cytoplasm Teratocarcinoma, Testis, alternative products: Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events, catalytic activity: A phosphoprotein + H(2)O = a protein + phosphata-confactor Binds 1 Fe(3+) in ope per subunit, confactor Binds 1 Fine subunit may have a role in the calmodulin activation of calcineurin, similarity. Belongs to the PPP phosphatase family, Ps-2 B subfamily, subunit: Composed of two components (A and B), the A component is the catalytic subunit and the B component confers calcium sensitivity. Background Background	Isotype	IgG
Gene Name PP93CB CALNA2 CALNB CNA2 Serine/threonine-protein phosphatase 2B catalytic subunit beta isoform (EC 3:1.3:16) (CAM-PRP catalytic subunit) (Calmodulin-dependent calcineurin A subunit beta isoform) Synthesized peptide derived from part region of human protein Specificity PP2BB Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 57kD Cell Pathway Cytoplasm. Teratocarcinoma,Testis, alternative products: Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events, catalytic activity: A phosphoprotein + H(2)O = a protein + phosphatase for calcineurin. similarity:Belongs to the PPP phosphatase family, subunit. Composed of two components (A and B), the A component is the catalytic subunit and the B component (A and B), the A component is the catalytic subunit and the B component is form alternative splicing events, catalytic activity: A phosphoprotein + H(2)O = a protein phosphatase family. Ps-2B subfamily, subunit:Composed of two components (A and B), the A component is the catalytic subunit and the B component is formal ternative splicing events, catalytic activity: A phosphoprotein + H(2)O = a protein + poducts: Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events, catalytic activity: A phosphoprotein + H(2)O = a protein +	Reactivity	Human;Rat;Mouse
Protein Name Serine/threonine-protein phosphatase 2B catalytic subunit beta isoform (EC 3.1.3.16) (CAMPRP catalytic subunit) (Calmodulin-dependent calcineurin A subunit beta isoform) Immunogen Synthesized peptide derived from part region of human protein PP2BB Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms Observed Band 57kD Cell Pathway Cytoplasm. Teratocarcinoma, Testis, Teratocarcinoma, Testis, Teratocarcinoma alternative products: Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events, catalytic activity-A phosphoprotein + H(2)O = a protein + phosphatase fools are collected. Calcineurin Similarity, Belongs to the PPP phosphatase family, subunit. Grom alternative splicing events, catalytic activity-A phosphoprotein + H(2)O = a protein + component is the catalytic subunit and the B component is far and B), the A component is the catalytic subunit and the B component is far and B), the A component is the catalytic subunit and the B component is far and B) is forms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events, catalytic activity-A phosphoprotein + H(2)O = a protein +	Applications	WB
3.1.3.16) (CAM-PRP catalytic subunit) (Calmodulin-dependent calcineurin A subunit beta isoform) Synthesized peptide derived from part region of human protein Specificity PP2BB Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 57kD Cell Pathway Cytoplasm. Teratocarcinoma, Testis, alternative products: Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events, catalylito activity: A phosphoprotein + H(2)O = a protein + phosphata, cofactor: Binds 1 Fe(4) ion per subunit, cofactor: Binds 1 Fe(4) ion per subunit, cofactor: Binds 1 Teratocarcinoments (A and B), the A component is the catalytic subunit and the B components (A and B), the A component is the catalytic subunit and the B component confers calcium sensitivity. alternative products: Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative products calcium sensitivity. alternative products: Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events, catalytic activity: A phosphoprotein + H(2)O = a protein +	Gene Name	PPP3CB CALNA2 CALNB CNA2
PP2BB Monoclonal Antibody detects endogenous levels of protein.	Protein Name	3.1.3.16) (CAM-PRP catalytic subunit) (Calmodulin-dependent calcineurin A
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 57kD Cell Pathway Cytoplasm. Tissue Specificity Teratocarcinoma, Testis, alternative products: Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events, catalytic activity: A phosphoprotein + H(2)0 = a protein + phosphate. cofactor: Binds 1 Fe(3+) ion per subunit. cofactor: Binds 1 zinc ion per subunit, function: Calcium-dependent, calmodulin-stimulated protein phosphatase This subunit may have a role in the calmodulin activation of calcineurin, similarity: Belongs to the PPP phosphatase family. PP-2B subfamily, subunit: Composed of two components (A and B), the A component is the catalytic subunit and the B component confers calcium sensitivity. alternative products: Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events, catalytic activity: A phosphoprotein + H(2)0 = a protein +	Immunogen	Synthesized peptide derived from part region of human protein
Monoclonal, Mouse, IgG	Specificity	PP2BB Monoclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. WB 1:500-1:2000 Concentration 1 mg/ml ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 57kD Cell Pathway Cytoplasm. Tissue Specificity Teratocarcinoma, Testis, alternative products: Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events, catalytic activity: A phosphoprotein + H(2)O = a protein + phosphatae, cofactor:Binds 1 Fe(3+) ion per subunit., cofactor:Binds 1 zinc ion per subunit. function: Calcineurin. A phosphatae family. PP-2B subfamily., similarity:Belongs to the PPP phosphataes family, similarity:Belongs to the PPP phosphataes family. subunit:Composed of two components (A and B), the A component is the catalytic subunit and the B component confers calcium sensitivity., alternative products:Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events, catalytic activity:A phosphoprotein + H(2)O = a protein + a catalytic subunit and the B component confers calcium sensitivity.,	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 57kD Cell Pathway Cytoplasm. Teratocarcinoma,Testis, alternative products:Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events, catalytic activity: A phosphoprotein + H(2)O = a protein + phosphate, cofactor:Binds 1 Fe(3+) ion per subunit, cofactor:Binds 1 zinc ion per subunit, function:Calcium-dependent, calmodulin-stimulated protein phosphatase This subunit may have a role in the calmodulin activation of calcineurin. similarity:Belongs to the PPP phosphatase family, similarity:Belongs to the PPP phosphatase family phosphatase family, subunit:Composed of two components (A and B), the A component is the catalytic subunit and the B component confers calcium sensitivity., alternative products:Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events, catalytic activity:A phosphoprotein + H(2)O = a protein +	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 57kD Cell Pathway Cytoplasm . Tissue Specificity Teratocarcinoma,Testis, alternative products:Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events, catalytic activity:A phosphoprotein + H(2)O = a protein + phosphate.,cofactor:Binds 1 Fe(3+) ion per subunit.,cofactor:Binds 1 zinc ion per subunit.,function:Calcium-dependent, calmodulin-stimulated protein phosphatase This subunit may have a role in the calmodulin-stimulated protein phosphatase This subunit may have a role in the calmodulin activation of calcineurin.,similarity:Belongs to the PPP phosphatase family. PP-2B subfamily.,subunit:Composed of two components (A and B), the A component is the catalytic subunit and the B component confers calcium sensitivity., alternative products:Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events, catalytic activity:A phosphoprotein + H(2)O = a protein +	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 57kD Cell Pathway Cytoplasm . Tissue Specificity Teratocarcinoma,Testis, alternative products:Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events,catalytic activity:A phosphoprotein + H(2)O = a protein + phosphate, cofactor:Binds 1 Fe(3+) ion per subunit, cofactor:Binds 1 zinc ion per subunit, function:Calcium-dependent, calmodulin-stimulated protein phosphatase This subunit may have a role in the calmodulin activation of calcineurin, similarity:Belongs to the PPP phosphatase family, subunit:Composed of two components (A and B), the A component is the catalytic subunit and the B component confers calcium sensitivity., Background alternative products:Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events, catalytic activity:A phosphoprotein + H(2)O = a protein +	Dilution	WB 1:500-1:2000
Synonyms Observed Band 57kD Cell Pathway Cytoplasm . Tissue Specificity Teratocarcinoma, Testis, alternative products: Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events, catalytic activity: A phosphoprotein + H(2)O = a protein + phosphate, cofactor: Binds 1 Fe(3+) ion per subunit, cofactor: Binds 1 zinc ion per subunit, function: Calcium-dependent, calmodulin-stimulated protein phosphatase This subunit may have a role in the calmodulin activation of calcineurin., similarity: Belongs to the PPP phosphatase family, similarity: Belongs to the PPP phosphatase family, subunit: Composed of two components (A and B), the A component is the catalytic subunit and the B component confers calcium sensitivity. Background alternative products: Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events, catalytic activity: A phosphoprotein + H(2)O = a protein +	Concentration	1 mg/ml
Observed Band 57kD Cell Pathway Cytoplasm . Tissue Specificity Teratocarcinoma, Testis, alternative products: Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events, catalytic activity: A phosphoprotein + H(2)O = a protein + phosphate., cofactor: Binds 1 Fe(3+) ion per subunit., cofactor: Binds 1 zinc ion per subunit., function: Calcium-dependent, calmodulin-stimulated protein phosphatase This subunit may have a role in the calmodulin-activation of calcineurin., similarity: Belongs to the PPP phosphatase family., similarity: Belongs to the PPP phosphatase family. PP-2B subfamily., subunit: Composed of two components (A and B), the A component is the catalytic subunit and the B component confers calcium sensitivity., alternative products: Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events, catalytic activity: A phosphoprotein + H(2)O = a protein +	Purity	≥90%
Observed Band Cell Pathway Cytoplasm . Tissue Specificity Teratocarcinoma, Testis, alternative products: Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events, catalytic activity: A phosphoprotein + H(2)O = a protein + phosphate., cofactor: Binds 1 Fe(3+) ion per subunit., cofactor: Binds 1 zinc ion per subunit., function: Calcium-dependent, calmodulin-stimulated protein phosphatase This subunit may have a role in the calmodulin activation of calcineurin., similarity: Belongs to the PPP phosphatase family. PP-2B subfamily., subunit: Composed of two components (A and B), the A component is the catalytic subunit and the B component confers calcium sensitivity. Background alternative products: Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events, catalytic activity: A phosphoprotein + H(2)O = a protein +	Storage Stability	-20°C/1 year
Cell Pathway Cytoplasm . Tissue Specificity Teratocarcinoma, Testis, alternative products: Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events, catalytic activity: A phosphoprotein + H(2)O = a protein + phosphate., cofactor: Binds 1 Fe(3+) ion per subunit., cofactor: Binds 1 zinc ion per subunit., function: Calcium-dependent, calmodulin-stimulated protein phosphatase This subunit may have a role in the calmodulin activation of calcineurin., similarity: Belongs to the PPP phosphatase family., similarity: Belongs to the PPP phosphatase family. PP-2B subfamily., subunit: Composed of two components (A and B), the A component is the catalytic subunit and the B component confers calcium sensitivity., alternative products: Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events, catalytic activity: A phosphoprotein + H(2)O = a protein +	Synonyms	
Tissue Specificity Teratocarcinoma, Testis, alternative products: Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events, catalytic activity: A phosphoprotein + H(2)O = a protein + phosphate., cofactor: Binds 1 Fe(3+) ion per subunit., cofactor: Binds 1 zinc ion per subunit., function: Calcium-dependent, calmodulin-stimulated protein phosphatase This subunit may have a role in the calmodulin activation of calcineurin., similarity: Belongs to the PPP phosphatase family., similarity: Belongs to the PPP phosphatase family. PP-2B subfamily., subunit: Composed of two components (A and B), the A component is the catalytic subunit and the B component confers calcium sensitivity., alternative products: Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events, catalytic activity: A phosphoprotein + H(2)O = a protein +	Observed Band	57kD
alternative products:Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events,catalytic activity:A phosphoprotein + H(2)O = a protein + phosphate.,cofactor:Binds 1 Fe(3+) ion per subunit.,cofactor:Binds 1 zinc ion per subunit.,function:Calcium-dependent, calmodulin-stimulated protein phosphatase This subunit may have a role in the calmodulin activation of calcineurin.,similarity:Belongs to the PPP phosphatase family.,similarity:Belongs to the PPP phosphatase family. PP-2B subfamily.,subunit:Composed of two components (A and B), the A component is the catalytic subunit and the B component confers calcium sensitivity., Background alternative products:Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events,catalytic activity:A phosphoprotein + H(2)O = a protein +	Cell Pathway	Cytoplasm .
consists of at least two isoenzymes that may result from alternative splicing events, catalytic activity: A phosphoprotein + H(2)O = a protein + phosphate., cofactor: Binds 1 Fe(3+) ion per subunit., cofactor: Binds 1 zinc ion per subunit., function: Calcium-dependent, calmodulin-stimulated protein phosphatase This subunit may have a role in the calmodulin activation of calcineurin., similarity: Belongs to the PPP phosphatase family., similarity: Belongs to the PPP phosphatase family. PP-2B subfamily., subunit: Composed of two components (A and B), the A component is the catalytic subunit and the B component confers calcium sensitivity., Background alternative products: Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events, catalytic activity: A phosphoprotein + H(2)O = a protein +	Tissue Specificity	Teratocarcinoma,Testis,
consists of at least two isoenzymes that may result from alternative splicing events catalytic activity:A phosphoprotein + H(2)O = a protein +	Function	events,catalytic activity:A phosphoprotein + H(2)O = a protein + phosphate.,cofactor:Binds 1 Fe(3+) ion per subunit.,cofactor:Binds 1 zinc ion per subunit.,function:Calcium-dependent, calmodulin-stimulated protein phosphatase This subunit may have a role in the calmodulin activation of calcineurin.,similarity:Belongs to the PPP phosphatase family.,similarity:Belongs to the PPP phosphatase family. PP-2B subfamily.,subunit:Composed of two components (A and B), the A component is the catalytic subunit and the B
	Background	alternative products:Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events,catalytic activity:A phosphoprotein + H(2)O = a protein + phosphate.,cofactor:Binds 1 Fe(3+) ion per subunit.,cofactor:Binds 1 zinc ion per



UpingBio technology Co.,Ltd





subunit.,function:Calcium-dependent, calmodulin-stimulated protein phosphatase. This subunit may have a role in the calmodulin activation of calcineurin.,similarity:Belongs to the PPP phosphatase family.,similarity:Belongs to the PPP phosphatase family. PP-2B subfamily.,subunit:Composed of two components (A and B), the A component is the catalytic subunit and the B component confers calcium sensitivity.,

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images