





LIRA1 Monoclonal Antibody

Catalog No	YP-mAb-07286
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	LILRA1 LIR6
Protein Name	Leukocyte immunoglobulin-like receptor subfamily A member 1 (CD85 antigen-like family member I) (Leukocyte immunoglobulin-like receptor 6) (LIR-6) (CD antigen CD85i)
Immunogen	Synthesized peptide derived from human protein . at AA range: Internal
Specificity	LIRA1 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
	>000/
Purity	≥90%
Storage Stability	-20°C/1 year
Storage Stability	
Storage Stability Synonyms	-20°C/1 year
Storage Stability Synonyms Observed Band	-20°C/1 year 53kD
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year 53kD Membrane; Single-pass type I membrane protein.
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year 53kD Membrane; Single-pass type I membrane protein. Detected in monocytes and B-cells. function:May act as receptor for class I MHC antigens.,similarity:Contains 4 Ig-like C2-type (immunoglobulin-like) domains.,tissue specificity:Detected in monocytes
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	-20°C/1 year 53kD Membrane; Single-pass type I membrane protein. Detected in monocytes and B-cells. function:May act as receptor for class I MHC antigens.,similarity:Contains 4 Ig-like C2-type (immunoglobulin-like) domains.,tissue specificity:Detected in monocytes and B-cells., This gene encodes an activating member of the leukocyte immunoglobulin-like receptor (LIR) family, which is found in a gene cluster at chromosomal region 19q13.4. The encoded protein is predominantly expressed in B cells, interacts with major histocompatibility complex class I ligands, and contributes to the regulation of immune responses. Alternative splicing results in multiple transcript



UpingBio technology Co.,Ltd





Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images