

THNS2 Monoclonal Antibody

Catalog No YP-mAb-07291 Isotype IgG Reactivity Human;Mouse;Rat Applications WB Gene Name THNSL2 Protein Name Threonine synthase-like 2 (TSH2) (EC 4.2.3) (Secreted osteoclastogenic factor of activated T-cells) (SOFAT) Immunogen Synthesized peptide derived from human protein . at AA range: 91-140 Specificity THNS2 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 53kD Cell Pathway [Isoform SOFAT]: Secreted. Secreted by activated T-cells via a calcineurin-independent pathway. Tissue Specificity Brain, Colon, Ovarian carcinoma, Skin, Tongue, Function Gofactor: Pyridoxal phosphate, function: Acts as a catabolic phospho-hyase on both gamma- and beta-phosphorylated substrates	Isotype Reactivity Applications Gene Name Protein Name	IgG Human;Mouse;Rat WB THNSL2 Threonine synthase-like 2 (TSH2) (EC 4.2.3) (Secreted osteoclastogenic factor of activated T-cells) (SOFAT) Synthesized peptide derived from human protein . at AA range: 91-140
Reactivity Human;Mouse;Rat Applications WB Gene Name THNSL2 Protein Name Threonine synthase-like 2 (TSH2) (EC 4.2.3) (Secreted osteoclastogenic factor of activated T-cells) (SOFAT) Immunogen Synthesized peptide derived from human protein . at AA range: 91-140 Specificity THNS2 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 53kD Cell Pathway [Isoform SOFAT]: Secreted, Secreted by activated T-cells via a calcineurin-independent pathway. Tissue Specificity Brain, Colon, Ovarian carcinoma, Skin, Tongue, Function cofactor: Pyridoxal phosphate, function: Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-kefobutyrate, ammonia and phosphate., similarity: Belongs to the serine/fthreonine and beta-phosphor-homoserine to a-ketobutyrate, phosphosphose, and ammonia. This protein also has phospho-lyase a activity on both gamma and beta-phosphorylated substrates. In mouse a alternatively spliced form of this protein has been shown to act as a volvokine and activity on both gamma and beta-phosphorylated substrates. In mouse an alternatively spliced form of this protein has been shown to act as a volvokine and activity on both gamma and beta-phosphorylated substrates. In mouse an alternatively spliced form of this protein has been shown to act as a volvokine and activity on both gamma and beta-phosphorylated substrates. In mouse an alternatively spliced form of this protein has been shown to act as a volvokine and calvidiven between the protein and the protein also has phosphorylated substrates.	Reactivity Applications Gene Name Protein Name	Human;Mouse;Rat WB THNSL2 Threonine synthase-like 2 (TSH2) (EC 4.2.3) (Secreted osteoclastogenic factor of activated T-cells) (SOFAT) Synthesized peptide derived from human protein . at AA range: 91-140
Applications WB Gene Name THNSL2 Protein Name Threonine synthase-like 2 (TSH2) (EC 4.2.3) (Secreted osteoclastogenic factor of activated T-cells) (SOFAT) (EC 4.2.3) (Secreted osteoclastogenic factor of activated T-cells) (SOFAT) (EC 4.2.3) (Secreted osteoclastogenic factor of activated T-cells) (SOFAT) (SOFAT) Immunogen Synthesized peptide derived from human protein . at AA range: 91-140 Specificity THNS2 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 53kD Cell Pathway [Isoform SOFAT]: Secreted. Secreted by activated T-cells via a calcineurin-independent pathway. Tissue Specificity Brain, Colon, Ovarian carcinoma, Skin, Tongue, Function cactor: Pyridoxal phosphate, function: Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-kefobutyrate, ammonia and phosphate., similarity: Belongs to the serine/threonine dehydratase familly. Background This gene encodes a threonine synthase-like protein. A similar enzyme in mouse can catalyze the degradation of O-phospho-homoserine to a-ketobutyrate, phosphate, and ammonia. This protein also has phospho-plase activity on both gamma and beta phosphorylated substrates. In mouse an alternatively spliced form of this protein has been shown to act as a cytokine and can induce the	Applications Gene Name Protein Name	WB THNSL2 Threonine synthase-like 2 (TSH2) (EC 4.2.3) (Secreted osteoclastogenic factor of activated T-cells) (SOFAT) Synthesized peptide derived from human protein . at AA range: 91-140
Gene Name THNSL2 Protein Name Threonine synthase-like 2 (TSH2) (EC 4.2.3) (Secreted osteoclastogenic factor of activated T-cells) (SOFAT) Immunogen Synthesized peptide derived from human protein . at AA range: 91-140 Specificity THNS2 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 53kD Cell Pathway [Isoform SOFAT]: Secreted. Secreted by activated T-cells via a calcineurin-independent pathway. Tissue Specificity Brain, Colon, Ovarian carcinoma, Skin, Tongue, Cofactor: Pyridoxal phosphorylated substrates a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and phosphate., similarity: Belongs to the serine/threonine dehydratase familly. Background This gene encodes a threonine synthase-like protein. A similar enzyme in mouse can catalyze the degradation of O-phospho-homoserine to a-ketobutyrate, phosphate, and ammonia. This protein also has phospho-lyase activity on both gamma and beta phosphorylated substrates. In mouse an alternatively spliced form of this protein has been shown to act as a cytokine and can induce the	Gene Name Protein Name	THNSL2 Threonine synthase-like 2 (TSH2) (EC 4.2.3) (Secreted osteoclastogenic factor of activated T-cells) (SOFAT) Synthesized peptide derived from human protein . at AA range: 91-140
Protein Name Threonine synthase-like 2 (TSH2) (EC 4.2.3) (Secreted osteoclastogenic factor of activated T-cells) (SOFAT) Immunogen Synthesized peptide derived from human protein . at AA range: 91-140 Specificity THNS2 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 53kD Cell Pathway [Isoform SOFAT]: Secreted. Secreted by activated T-cells via a calcineurin-independent pathway. Tissue Specificity Brain, Colon, Ovarian carcinoma, Skin, Tongue, cofactor: Pyridoxal phosphate., function: Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and phosphate., similarity: Belongs to the serine/threonine dehydratase family. Background This gene encodes a threonine synthase-like protein. A similar enzyme in mouse can catalyze the degradation of O-phospho-homoserine to a-ketobutyrate, phosphate, and ammonia. This protein also has phosphor-lyase activity on both gamma and beta phosphorylated substrates. In mouse an alternatively spliced form of this protein has been shown to act as a cytokine and can induce the	Protein Name	Threonine synthase-like 2 (TSH2) (EC 4.2.3) (Secreted osteoclastogenic factor of activated T-cells) (SOFAT) Synthesized peptide derived from human protein . at AA range: 91-140
of activated T-cells) (SOFAT) Immunogen Synthesized peptide derived from human protein . at AA range: 91-140 Specificity THNS2 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 53kD Cell Pathway [Isoform SOFAT]: Secreted. Secreted by activated T-cells via a calcineurin-independent pathway. Tissue Specificity Brain, Colon, Ovarian carcinoma, Skin, Tongue, Function cofactor: Pyridoxal phosphate, function: Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PTnr) to alpha-ketobutyrate, ammonia and phosphate, similarity: Belongs to the serine/threonine dehydratase family. Background This gene encodes a threonine synthase-like protein. A similar enzyme in mouse can catalyze the degradation of O-phospho-homoserine to a-ketobutyrate, phosphate, ammonia and phosphate. In mouse an alternatively spliced form of this protein also has phospho-lyase activity on both gamma and beta phosphorylated substrates. In mouse an alternatively spliced form of this protein has been shown to act as a cytokine and can induce the		of activated T-cells) (SOFAT) Synthesized peptide derived from human protein . at AA range: 91-140
Specificity THNS2 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Cell Pathway Cell Pathway [Isoform SOFAT]: Secreted. Secreted by activated T-cells via a calcineurin-independent pathway. Tissue Specificity Brain, Colon, Ovarian carcinoma, Skin, Tongue, Function cofactor: Pyridoxal phosphate., function: Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PTThr) to alpha-ketobutyrate, ammonia and phosphate., similarity: Belongs to the serine/fithreonine dehydratase family. Background This gene encodes a threonine synthase-like protein. A similar enzyme in mouse can catalyze the degradation of O-phospho-homoserine to a-ketobutyrate, phosphate, and ammonia. This protein also has phospho-lyase activity on both gamma and beta phosphorylated substrates. In mouse an alternatively spliced form of this protein has been shown to act as a cytokine and can induce the	Immunogen	
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 53kD Cell Pathway [Isoform SOFAT]: Secreted. Secreted by activated T-cells via a calcineurin-independent pathway. Tissue Specificity Brain, Colon, Ovarian carcinoma, Skin, Tongue, Function cofactor: Pyridoxal phosphate., function: Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and phosphate., similarity: Belongs to the serine/threonine dehydratase family. This gene encodes a threonine synthase-like protein. A similar enzyme in mouse can catalyze the degradation of O-phospho-homoserine to a-ketobutyrate, phosphate, and ammonia. This protein also has phospho-lyase activity on both gamma and beta phosphorylated substrates. In mouse an alternatively spliced form of this protein has been shown to act as a cytokine and can induce the		TINOON I IN OUR IN THE STATE OF
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 53kD Cell Pathway [Isoform SOFAT]: Secreted. Secreted by activated T-cells via a calcineurin-independent pathway. Tissue Specificity Brain,Colon,Ovarian carcinoma,Skin,Tongue, Function cofactor:Pyridoxal phosphate.,function:Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and phosphate.,similarity:Belongs to the serine/threonine dehydratase family., Background This gene encodes a threonine synthase-like protein. A similar enzyme in mouse can catalyze the degradation of O-phospho-homoserine to a-ketobutyrate, phosphate, and ammonia. This protein also has phospho-lyase activity on both gamma and beta phosphorylated substrates. In mouse can alternatively spliced form of this protein has been shown to act as a cytokine and can induce the	Specificity	I HINSZ Monocional Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 53kD Cell Pathway [Isoform SOFAT]: Secreted. Secreted by activated T-cells via a calcineurin-independent pathway. Tissue Specificity Brain, Colon, Ovarian carcinoma, Skin, Tongue, Function cofactor: Pyridoxal phosphate., function: Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and phosphate., similar renzyme in mouse can catalyze the degradation of O-phospho-homoserine to a-ketobutyrate, phosphate, and ammonia. This protein also has phospho-lyase activity on both gamma and beta phosphorylated substrates. In mouse an alternatively spliced form of this protein has been shown to act as a cytokine and can induce the	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 53kD Cell Pathway [Isoform SOFAT]: Secreted. Secreted by activated T-cells via a calcineurin-independent pathway. Tissue Specificity Brain,Colon,Ovarian carcinoma,Skin,Tongue, Function cofactor:Pyridoxal phosphate.,function:Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and phosphate.,similarity:Belongs to the serine/threonine dehydratase family. Background This gene encodes a threonine synthase-like protein. A similar enzyme in mouse can catalyze the degradation of O-phospho-homoserine to a-ketobutyrate, phosphate, and ammonia. This protein also has phospho-lyase activity on both gamma and beta phosphorylated substrates. In mouse an alternatively spliced form of this protein has been shown to act as a cytokine and can induce the	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 53kD Cell Pathway [Isoform SOFAT]: Secreted. Secreted by activated T-cells via a calcineurin-independent pathway. Tissue Specificity Brain,Colon,Ovarian carcinoma,Skin,Tongue, Function cofactor:Pyridoxal phosphate.,function:Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and phosphate.,similarity:Belongs to the serine/threonine dehydratase family. Background This gene encodes a threonine synthase-like protein. A similar enzyme in mouse can catalyze the degradation of O-phospho-homoserine to a-ketobutyrate, phosphate, and ammonia. This protein also has phospho-lyase activity on both gamma and beta phosphorylated substrates. In mouse an alternatively spliced form of this protein has been shown to act as a cytokine and can induce the	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 53kD Cell Pathway [Isoform SOFAT]: Secreted. Secreted by activated T-cells via a calcineurin-independent pathway. Tissue Specificity Brain,Colon,Ovarian carcinoma,Skin,Tongue, Function cofactor:Pyridoxal phosphate,,function:Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and phosphate,,similarity:Belongs to the serine/threonine dehydratase family., Background This gene encodes a threonine synthase-like protein. A similar enzyme in mouse can catalyze the degradation of O-phospho-homoserine to a-ketobutyrate, phosphate, and ammonia. This protein also has phospho-lyase activity on both gamma and beta phosphorylated substrates. In mouse an alternatively spliced form of this protein has been shown to act as a cytokine and can induce the	Dilution	WB 1:500-1:2000
Storage Stability -20°C/1 year Synonyms Observed Band 53kD Cell Pathway [Isoform SOFAT]: Secreted. Secreted by activated T-cells via a calcineurin-independent pathway. Tissue Specificity Brain,Colon,Ovarian carcinoma,Skin,Tongue, Function cofactor:Pyridoxal phosphate.,function:Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and phosphate.,similarity:Belongs to the serine/threonine dehydratase family., Background This gene encodes a threonine synthase-like protein. A similar enzyme in mouse can catalyze the degradation of O-phospho-homoserine to a-ketobutyrate, phosphate, and ammonia. This protein also has phospho-lyase activity on both gamma and beta phosphorylated substrates. In mouse an alternatively spliced form of this protein has been shown to act as a cytokine and can induce the	Concentration	1 mg/ml
Synonyms Observed Band 53kD Cell Pathway [Isoform SOFAT]: Secreted. Secreted by activated T-cells via a calcineurin-independent pathway. Tissue Specificity Brain, Colon, Ovarian carcinoma, Skin, Tongue, Function cofactor: Pyridoxal phosphate., function: Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and phosphate., similarity: Belongs to the serine/threonine dehydratase family., Background This gene encodes a threonine synthase-like protein. A similar enzyme in mouse can catalyze the degradation of O-phospho-homoserine to a-ketobutyrate, phosphate, and ammonia. This protein also has phospho-lyase activity on both gamma and beta phosphorylated substrates. In mouse an alternatively spliced form of this protein has been shown to act as a cytokine and can induce the	Purity	≥90%
Observed Band Cell Pathway [Isoform SOFAT]: Secreted. Secreted by activated T-cells via a calcineurin-independent pathway. Tissue Specificity Brain,Colon,Ovarian carcinoma,Skin,Tongue, cofactor:Pyridoxal phosphate.,function:Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and phosphate.,similarity:Belongs to the serine/threonine dehydratase family., Background This gene encodes a threonine synthase-like protein. A similar enzyme in mouse can catalyze the degradation of O-phospho-homoserine to a-ketobutyrate, phosphate, and ammonia. This protein also has phospho-lyase activity on both gamma and beta phosphorylated substrates. In mouse an alternatively spliced form of this protein has been shown to act as a cytokine and can induce the	Storage Stability	-20°C/1 year
Cell Pathway [Isoform SOFAT]: Secreted. Secreted by activated T-cells via a calcineurin-independent pathway. Tissue Specificity Brain,Colon,Ovarian carcinoma,Skin,Tongue, Cofactor:Pyridoxal phosphate.,function:Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and phosphate.,similarity:Belongs to the serine/threonine dehydratase family., Background This gene encodes a threonine synthase-like protein. A similar enzyme in mouse can catalyze the degradation of O-phospho-homoserine to a-ketobutyrate, phosphate, and ammonia. This protein also has phospho-lyase activity on both gamma and beta phosphorylated substrates. In mouse an alternatively spliced form of this protein has been shown to act as a cytokine and can induce the	Synonyms	
Tissue Specificity Brain,Colon,Ovarian carcinoma,Skin,Tongue, Cofactor:Pyridoxal phosphate.,function:Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and phosphate.,similarity:Belongs to the serine/threonine dehydratase family., Background This gene encodes a threonine synthase-like protein. A similar enzyme in mouse can catalyze the degradation of O-phospho-homoserine to a-ketobutyrate, phosphate, and ammonia. This protein also has phospho-lyase activity on both gamma and beta phosphorylated substrates. In mouse an alternatively spliced form of this protein has been shown to act as a cytokine and can induce the	Observed Band	53kD
Function cofactor:Pyridoxal phosphate.,function:Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and phosphate.,similarity:Belongs to the serine/threonine dehydratase family., This gene encodes a threonine synthase-like protein. A similar enzyme in mouse can catalyze the degradation of O-phospho-homoserine to a-ketobutyrate, phosphate, and ammonia. This protein also has phospho-lyase activity on both gamma and beta phosphorylated substrates. In mouse an alternatively spliced form of this protein has been shown to act as a cytokine and can induce the	Cell Pathway	
gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and phosphate.,similarity:Belongs to the serine/threonine dehydratase family., Background This gene encodes a threonine synthase-like protein. A similar enzyme in mouse can catalyze the degradation of O-phospho-homoserine to a-ketobutyrate, phosphate, and ammonia. This protein also has phospho-lyase activity on both gamma and beta phosphorylated substrates. In mouse an alternatively spliced form of this protein has been shown to act as a cytokine and can induce the	Tissue Specificity	Brain, Colon, Ovarian carcinoma, Skin, Tongue,
can catalyze the degradation of O-phospho-homoserine to a-ketobutyrate, phosphate, and ammonia. This protein also has phospho-lyase activity on both gamma and beta phosphorylated substrates. In mouse an alternatively spliced form of this protein has been shown to act as a cytokine and can induce the	Function	gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and phosphate similarity:Belongs to the
results in multiple transcript variants. [provided by RefSeq, Sep 2011],	Background	can catalyze the degradation of O-phospho-homoserine to a-ketobutyrate, phosphate, and ammonia. This protein also has phospho-lyase activity on both gamma and beta phosphorylated substrates. In mouse an alternatively spliced form of this protein has been shown to act as a cytokine and can induce the production of the inflammatory cytokine IL6 in osteoblasts. Alternate splicing
matters needing Avoid repeated freezing and thawing! attention		Avoid repeated freezing and thawing!



UpingBio technology Co.,Ltd





Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images