



CASS4 Monoclonal Antibody

Catalog No	YP-mAb-07661
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	CASS4 C20orf32 HEFL
Protein Name	Cas scaffolding protein family member 4 (HEF-like protein) (HEF1-EFS-p130Cas-like protein) (HEPL)
Immunogen	Synthesized peptide derived from part region of human protein AA range: 302-352
Specificity	CASS4 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	86kD
Cell Pathway	Cytoplasm, cytoskeleton . Cell junction, focal adhesion .
Tissue Specificity	Expressed abundantly in lung and spleen. Also highly expressed in ovarian and leukemia cell lines.
Function	alternative products:Experimental confirmation may be lacking for some isoforms,function:Possible docking protein which may play a role for tyrosine-kinase-based signaling related to cell adhesion. Regulates FAK activity, focal adhesion integrity, and cell spreading.,PTM:Phosphorylated on tyrosines by SRC.,similarity:Belongs to the CAS family.,similarity:Contains 1 SH3 domain.,subunit:Interacts (via SH3 domain) with focal adhesion kinase (via C-terminus).,tissue specificity:Expressed abundantly in lung and spleen. Also highly expressed in ovarian and leukemia cell lines.,
Background	alternative products:Experimental confirmation may be lacking for some isoforms,function:Possible docking protein which may play a role for tyrosine-kinase-based signaling related to cell adhesion. Regulates FAK activity, focal adhesion integrity, and cell spreading.,PTM:Phosphorylated on tyrosines by SRC.,similarity:Belongs to the CAS family.,similarity:Contains 1 SH3 domain.,subunit:Interacts (via SH3 domain) with focal adhesion kinase (via



C-terminus).,tissue specificity:Expressed abundantly in lung and spleen. Also highly expressed in ovarian and leukemia cell lines.,

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images