

Webslte: www.upingBio.com

## DDX5 Monoclonal Antibody

Catalog No	YP-mAb-07667
Isotype	lgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	DDX5 G17P1 HELR HLR1
Protein Name	Probable ATP-dependent RNA helicase DDX5 (EC 3.6.4.13) (DEAD box protein 5) (RNA helicase p68)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	DDX5 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Synonyms Observed Band	
	Nucleus . Nucleus, nucleolus . Cytoplasm . During the G0 phase, predominantly located in the nucleus. Cytoplasmic levels increase during the G1/S phase. During the M phase, located at the vicinity of the condensed chromosomes. At G1, localizes in the cytoplasm
Observed Band	located in the nucleus. Cytoplasmic levels increase during the G1/S phase. During the M phase, located at the vicinity of the condensed chromosomes. At
Observed Band Cell Pathway	located in the nucleus. Cytoplasmic levels increase during the G1/S phase. During the M phase, located at the vicinity of the condensed chromosomes. At G1, localizes in the cytoplasm.
Observed Band Cell Pathway Tissue Specificity	located in the nucleus. Cytoplasmic levels increase during the G1/S phase. During the M phase, located at the vicinity of the condensed chromosomes. At G1, localizes in the cytoplasm.
Observed Band Cell Pathway Tissue Specificity Function	located in the nucleus. Cytoplasmic levels increase during the G1/S phase. During the M phase, located at the vicinity of the condensed chromosomes. At G1, localizes in the cytoplasm.



S Tel: 400-999-8863 Emall: UpingBio@163.com

Webslte: www.upingBio.com

## **Products Images**