



SEPT4 Monoclonal Antibody

Catalog No	YP-mAb-07771
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	SEPT4 ARTS PNUTL2 SEP4 hucep-7
Protein Name	Septin-4 (Apoptosis-related protein in the TGF-beta signaling pathway) (ARTS) (Bradeion beta) (Brain protein H5) (CE5B3 beta) (Cell division control-related protein 2) (hCDCREL-2) (Cerebral protein 7)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	SEPT4 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	52kD
Cell Pathway	Cytoplasm . Cell projection, cilium, flagellum . Cytoplasmic vesicle, secretory vesicle . Cell projection, axon . Cell projection, dendrite . Perikaryon . Cell junction, synapse . In platelets, found in areas surrounding alpha-granules (PubMed:15116257). Found in the sperm annulus, a fibrous ring structure connecting the midpiece and the principal piece of the sperm flagellum (PubMed:25588830). Expressed and colocalized with SLC6A3 and SNCA in axon terminals, especially at the varicosities (By similarity). .; [Isoform ARTS]: Mitochondrion . Nucleus . While predominantly localized in the mitochondria under resting conditions, translocates into the nucleus after TGF-beta treatment and apoptosis induction. .
Tissue Specificity	Widely expressed in adult and fetal tissues with highest expression in adult brain (at protein level), heart, liver and adrenal gland and fetal heart, kidney, liver and lung. Expressed in presynaptic terminals of dopaminergic neurons projecting from the substantia nigra pars compacta to the striatum (at protein level) (PubMed:17296554). Expressed in axonal varicosities in dopaminergic nerve terminals (at protein level) (PubMed:17296554). Expressed in the putamen and in the adjacent cerebral cortex (at protein level) (PubMed:17296554). Expressed in colonic crypts (at protein level) (PubMed:30389919). Also expressed in colorectal cancers and malignant melanomas. Expressed in platelets. .; [Isoform ARTS]:



Highly expressed in the brain and heart.

Function

function:Involved in cytokinesis .,similarity:Belongs to the septin family.,subcellular location:Only observed in tumor tissue cells.,subunit:May assemble into a multicomponent structure.,tissue specificity:Expressed ubiquitously in adult and fetal tissues with particularly high expression in adult heart, liver and adrenal gland and fetal heart, kidney, liver and lung. Also expressed in colorectal cancer and malignant melanoma samples.,

Background

septin 4(SEPT4) Homo sapiens This gene is a member of the septin family of nucleotide binding proteins, originally described in yeast as cell division cycle regulatory proteins. Septins are highly conserved in yeast, Drosophila, and mouse, and appear to regulate cytoskeletal organization. Disruption of septin function disturbs cytokinesis and results in large multinucleate or polyploid cells. This gene is highly expressed in brain and heart. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. One of the isoforms (known as ARTS) is distinct; it is localized to the mitochondria, and has a role in apoptosis and cancer. [provided by RefSeq, Nov 2010],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images