





DIAP2 Monoclonal Antibody

Catalog No	YP-mAb-07773
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	DIAPH2 DIA
Protein Name	Protein diaphanous homolog 2 (Diaphanous-related formin-2) (DRF2)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	DIAP2 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Synonyms Observed Band	
-	[Isoform 3]: Cytoplasm, cytosol. Early endosome. Isoform 3 is cytosolic but when coexpressed with RHOD, the 2 proteins colocalize to early endosomes.
Observed Band	[Isoform 3]: Cytoplasm, cytosol. Early endosome. Isoform 3 is cytosolic but when coexpressed with RHOD, the 2 proteins colocalize to early endosomes. Expressed in testis, ovary, small intestine, prostate, lung, liver, kidney and leukocytes.
Observed Band Cell Pathway	Expressed in testis, ovary, small intestine, prostate, lung, liver, kidney and
Observed Band Cell Pathway Tissue Specificity	Expressed in testis, ovary, small intestine, prostate, lung, liver, kidney and
Observed Band Cell Pathway Tissue Specificity Function	Expressed in testis, ovary, small intestine, prostate, lung, liver, kidney and
Observed Band Cell Pathway Tissue Specificity Function Background matters needing	Expressed in testis, ovary, small intestine, prostate, lung, liver, kidney and leukocytes.



UpingBio technology Co.,Ltd





Products Images