



DSC3 Monoclonal Antibody

Catalog No	YP-mAb-07793
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	DSC3 CDHF3 DSC4
Protein Name	Desmocollin-3 (Cadherin family member 3) (Desmocollin-4) (HT-CP)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	DSC3 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	98kD
Cell Pathway	Cell membrane; Single-pass type I membrane protein. Cell junction, desmosome.
Tissue Specificity	Epidermis, buccal mucosa, esophagus and cervix.
Function	domain:Calcium may be bound by the cadherin-like repeats .,function:Component of intercellular desmosome junctions. Involved in the interaction of plaque proteins and intermediate filaments mediating cell-cell adhesion. May contribute to epidermal cell positioning (stratification) by mediating differential adhesiveness between cells that express different isoforms.,similarity:Contains 5 cadherin domains.,tissue specificity:Epidermis, buccal mucosa, esophagus and cervix.,
Background	The protein encoded by this gene is a calcium-dependent glycoprotein that is a member of the desmocollin subfamily of the cadherin superfamily. These desmosomal family members, along with the desmogleins, are found primarily in epithelial cells where they constitute the adhesive proteins of the desmosome cell-cell junction and are required for cell adhesion and desmosome formation. The desmosomal family members are arranged in two clusters on chromosome 18, occupying less than 650 kb combined. Mutations in this gene are a cause of hypotrichosis and recurrent skin vesicles disorder. The protein can act as an autoantigen in pemphigus diseases, and it is also considered to be a biomarker



for some cancers. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq, Apr 2014],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images