



E4F1 mouse mAb

Catalog No	YP-mAb-09242
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	E4F1 E4F
Protein Name	Transcription factor E4F1 (EC 6.3.2) (E4F transcription factor 1) (Putative E3 ubiquitin-protein ligase E4F1) (Transcription factor E4F) (p120E4F) (p50E4F)
Immunogen	Synthesized peptide derived from human E4F1
Specificity	This antibody detects endogenous levels of human E4F1
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
	•
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year Nucleus, nucleoplasm. Cytoplasm. A small fraction is detected in the cytoplasm. Excluded from the nucleolus where it is targeted upon CDKN2A overexpression.
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year Nucleus, nucleoplasm. Cytoplasm. A small fraction is detected in the cytoplasm. Excluded from the nucleolus where it is targeted upon CDKN2A overexpression. Localizes to the mitotic spindle during embryogenesis
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year Nucleus, nucleoplasm. Cytoplasm. A small fraction is detected in the cytoplasm. Excluded from the nucleolus where it is targeted upon CDKN2A overexpression. Localizes to the mitotic spindle during embryogenesis
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year Nucleus, nucleoplasm. Cytoplasm. A small fraction is detected in the cytoplasm. Excluded from the nucleolus where it is targeted upon CDKN2A overexpression. Localizes to the mitotic spindle during embryogenesis
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90% -20°C/1 year Nucleus, nucleoplasm. Cytoplasm. A small fraction is detected in the cytoplasm. Excluded from the nucleolus where it is targeted upon CDKN2A overexpression. Localizes to the mitotic spindle during embryogenesis Ubiquitously expressed.



UpingBio technology Co.,Ltd





Products Images