



NECD mouse mAb

Catalog No	YP-mAb-09431
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	NDN
Protein Name	Necdin
Immunogen	Synthesized peptide derived from human NECD AA range: 101-151
Specificity	This antibody detects endogenous levels of human NECD
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Perikaryon. Nucleus. Neural perikarya, translocates to the nucleus of postmitotic neurons and interacts with the nuclear matrix.
Tissue Specificity	Almost ubiquitous. Detected in fetal brain, lung, liver and kidney; in adult heart, brain, placenta, lung, liver, skeletal muscle, kidney, pancreas, spleen, thymus, prostate, testis, ovary, small intestine and colon. Not detected in peripheral blood leukocytes. In brain, restricted to post-mitotic neurons.
Function	
Background	
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images