







ECM2 mouse mAb

| Catalog No | YP-mAb-09632 |
|---|--|
| Isotype | IgG |
| Reactivity | Human;Mouse |
| Applications | WB |
| Gene Name | ECM2 |
| Protein Name | Extracellular matrix protein 2 (Matrix glycoprotein SC1/ECM2) |
| Immunogen | Synthesized peptide derived from human ECM2 AA range: 101-151 |
| Specificity | This antibody detects endogenous levels of human ECM2 |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-1:2000 |
| Concentration | 1 mg/ml |
| | · ··· ·g ····· |
| Purity | ≥90% |
| | - |
| Purity | ≥90% |
| Purity Storage Stability | ≥90% |
| Purity Storage Stability Synonyms | ≥90% |
| Purity Storage Stability Synonyms Observed Band | ≥90% -20°C/1 year |
| Purity Storage Stability Synonyms Observed Band Cell Pathway | ≥90% -20°C/1 year Secreted, extracellular space, extracellular matrix . Expressed predominantly in adipose tissue as well as female-specific organs |
| Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity | ≥90% -20°C/1 year Secreted, extracellular space, extracellular matrix . Expressed predominantly in adipose tissue as well as female-specific organs |
| Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function | ≥90% -20°C/1 year Secreted, extracellular space, extracellular matrix . Expressed predominantly in adipose tissue as well as female-specific organs |
| Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing | ≥90% -20°C/1 year Secreted, extracellular space, extracellular matrix . Expressed predominantly in adipose tissue as well as female-specific organs such as mammary gland, ovary, and uterus. |



UpingBio technology Co.,Ltd





Products Images