





MYRIP mouse mAb

| Catalog No | YP-mAb-09680 |
|---------------------------|---|
| Isotype | IgG |
| Reactivity | Human;Mouse;Rat |
| Applications | WB |
| Gene Name | MYRIP SLAC2C |
| Protein Name | Rab effector MyRIP (Exophilin-8) (Myosin-VIIa- and Rab-interacting protein) (Synaptotagmin-like protein lacking C2 domains C) (SlaC2-c) (Slp homolog lacking C2 domains c) |
| Immunogen | Synthesized peptide derived from human MYRIP AA range: 591-641 |
| Specificity | This antibody detects endogenous levels of human MYRIP |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-1:2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | |
| Cell Pathway | Cytoplasm . Cytoplasm, perinuclear region . Cytoplasmic vesicle, secretory vesicle . In presynaptic and postsynaptic areas in photoreceptor cells and in the basal microvilli of retinal pigment epithelium cells. Associated with melanosomes Colocalizes with actin filaments |
| Tissue Specificity | Detected in brain, skin, heart, adrenal medulla, pancreas, intestine, liver, kidney, muscle and testis. |
| Function | |
| Background | |
| matters needing attention | Avoid repeated freezing and thawing! |
| Usage suggestions | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |
| | |



UpingBio technology Co.,Ltd





Products Images