







TRAT1 mouse mAb

Catalog No	YP-mAb-09708
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	TRAT1 TCRIM HSPC062
Protein Name	T-cell receptor-associated transmembrane adapter 1 (T-cell receptor-interacting molecule) (TRIM) (pp29/30)
Immunogen	Synthesized peptide derived from human TRAT1 AA range: 56-106
Specificity	This antibody detects endogenous levels of human TRAT1
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	1 mg/ml ≥90%
	•
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year Cell membrane ; Single-pass type III membrane protein . Strongly expressed in thymus, and to a lesser extent in spleen, lymph node and peripheral blood lymphocytes. Present in T-cells and NK cells, but not B-cells (at
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year Cell membrane ; Single-pass type III membrane protein . Strongly expressed in thymus, and to a lesser extent in spleen, lymph node and peripheral blood lymphocytes. Present in T-cells and NK cells, but not B-cells (at
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year Cell membrane ; Single-pass type III membrane protein . Strongly expressed in thymus, and to a lesser extent in spleen, lymph node and peripheral blood lymphocytes. Present in T-cells and NK cells, but not B-cells (at
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90% -20°C/1 year Cell membrane ; Single-pass type III membrane protein . Strongly expressed in thymus, and to a lesser extent in spleen, lymph node and peripheral blood lymphocytes. Present in T-cells and NK cells, but not B-cells (at protein level).



UpingBio technology Co.,Ltd





Products Images