



UBQL1 mouse mAb

Catalog No	YP-mAb-10079
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	UBQLN1 DA41 PLIC1
Protein Name	Ubiquilin-1 (Protein linking IAP with cytoskeleton 1) (PLIC-1) (hPLIC-1)
Immunogen	Synthesized peptide derived from human UBQL1 AA range: 302-352
Specificity	This antibody detects endogenous levels of human UBQL1
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Cytoplasm . Nucleus . Endoplasmic reticulum . Cytoplasmic vesicle, autophagosome . Cell membrane . Detected in neuronal processes and at synapses (By similarity). Recruited to the ER during ER-associated protein degradation (ERAD) (PubMed:19822669). Isoform 1 and isoform 3 colocalize with PSEN1 in the cell membrane and in cytoplasmic juxtanuclear structures called aggresomes (PubMed:21143716). Colocalizes with ORAI1 and TICAM1 in the autophagosome (PubMed:23307288, PubMed:21695056). Colocalizes with EPS15 and HGS in ubiquitin-rich cytoplasmic aggregates that are not endocytic compartments and with EPS15 also in aggresomes (PubMed:16159959). .
Tissue Specificity	Brain (at protein level) (PubMed:18953672). Ubiquitous. Highly expressed throughout the brain; detected in neurons and in neuropathological lesions, such as neurofibrillary tangles and Lewy bodies. Highly expressed in heart, placenta, pancreas, lung, liver, skeletal muscle and kidney.
Function	
Background	

**matters needing attention**

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images