

ACAD8 mouse mAb

Catalog No	YP-mAb-10136
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	ACAD8 ARC42 IBD
Protein Name	Isobutyryl-CoA dehydrogenase, mitochondrial (EC 1.3.99) (Activator-recruited cofactor 42 kDa component) (ARC42) (Acyl-CoA dehydrogenase family member 8) (ACAD-8)
Immunogen	Synthesized peptide derived from human ACAD8 AA range: 326-376
Specificity	This antibody detects endogenous levels of human ACAD8
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
	≥90% -20°C/1 year
Purity	
Purity Storage Stability	
Purity Storage Stability Synonyms	
Purity Storage Stability Synonyms Observed Band	-20°C/1 year Mitochondrion .
Purity Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year Mitochondrion . Detected at comparable levels in all tissues examined (heart, lung, brain, skeletal
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year Mitochondrion . Detected at comparable levels in all tissues examined (heart, lung, brain, skeletal
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	-20°C/1 year Mitochondrion . Detected at comparable levels in all tissues examined (heart, lung, brain, skeletal
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	-20°C/1 year Mitochondrion . Detected at comparable levels in all tissues examined (heart, lung, brain, skeletal muscle, pancreas and placenta). Weakly expressed in liver and kidney.



UpingBio technology Co.,Ltd





Products Images