

## ISOC2 mouse mAb

| Catalog No  | YP-mAb-10225   |
|---|--|
| Isotype   | IgG  |
| Reactivity  | Human;Rat  |
| Applications  | WB   |
| Gene Name   | ISOC2  |
| Protein Name  | Isochorismatase domain-containing protein 2, mitochondrial   |
| Immunogen   | Synthesized peptide derived from human ISOC2 AA range: 47-97   |
| Specificity   | This antibody detects endogenous levels of human ISOC2   |
| Formulation   | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| Source  | Monoclonal, Mouse,IgG  |
| Purification  | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution  | WB 1:500-1:2000  |
| Concentration   | 1 mg/ml  |
| Purity  | ≥90%   |
| Storage Stability   | -20°C/1 year   |
|   |  |
| Synonyms  |  |
| Synonyms Observed Band  |  |
|   | Cytoplasm . Nucleus . Localizes to the nucleus in the presence of CDKN2A.  |
| Observed Band   | Cytoplasm . Nucleus . Localizes to the nucleus in the presence of CDKN2A.  |
| Observed Band Cell Pathway  | Cytoplasm . Nucleus . Localizes to the nucleus in the presence of CDKN2A.  |
| Observed Band Cell Pathway Tissue Specificity                     | Cytoplasm . Nucleus . Localizes to the nucleus in the presence of CDKN2A.  |
| Observed Band Cell Pathway Tissue Specificity Function            | Cytoplasm . Nucleus . Localizes to the nucleus in the presence of CDKN2A.  Avoid repeated freezing and thawing!      |
| Observed Band Cell Pathway Tissue Specificity Function Background |  |



## UpingBio technology Co.,Ltd





## **Products Images**