





## ZKSC3 mouse mAb

Catalog No	YP-mAb-10242
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	ZKSCAN3 ZFP47 ZNF306 ZNF309 ZSCAN13
Protein Name	Zinc finger protein with KRAB and SCAN domains 3 (Zinc finger and SCAN domain-containing protein 13) (Zinc finger protein 306) (Zinc finger protein 309) (Zinc finger protein 47 homolog) (Zf47) (Zfp-47
Immunogen	Synthesized peptide derived from human ZKSC3 AA range: 457-507
Specificity	This antibody detects endogenous levels of human ZKSC3
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,lgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year  Nucleus. Cytoplasm. Mainly localizes in the nucleus. Under starvation conditions translocates to the cytoplasm, allowing expression of target genes involved in
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year  Nucleus. Cytoplasm. Mainly localizes in the nucleus. Under starvation conditions translocates to the cytoplasm, allowing expression of target genes involved in
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year  Nucleus. Cytoplasm. Mainly localizes in the nucleus. Under starvation conditions translocates to the cytoplasm, allowing expression of target genes involved in
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year  Nucleus. Cytoplasm. Mainly localizes in the nucleus. Under starvation conditions translocates to the cytoplasm, allowing expression of target genes involved in
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90%  -20°C/1 year  Nucleus. Cytoplasm. Mainly localizes in the nucleus. Under starvation conditions translocates to the cytoplasm, allowing expression of target genes involved in autophagy and lysosome biogenesis/function.



## UpingBio technology Co.,Ltd





## **Products Images**