



# Myosin IIa (phospho-Ser1943) mouse mAb

<b>Catalog No</b>	YP-mAb-10384
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	WB
<b>Gene Name</b>	MYH9
<b>Protein Name</b>	Myosin IIa (Ser1943)
<b>Immunogen</b>	Synthesized phosho peptide around human Myosin IIa (Ser1943)
<b>Specificity</b>	This antibody detects endogenous levels of Human Mouse Rat Myosin IIa (phospho-Ser1943)
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-1:2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	Myosin-9 (Cellular myosin heavy chain, type A) (Myosin heavy chain 9) (Myosin heavy chain, non-muscle IIa) (Non-muscle myosin heavy chain A) (NMMHC-A) (Non-muscle myosin heavy chain IIa) (NMMHC II-a) (NMMHC-IIA)
<b>Observed Band</b>	100430DA
<b>Cell Pathway</b>	Cytoplasm, cytoskeleton . Cytoplasm, cell cortex . Cytoplasmic vesicle, secretory vesicle, Cortical granule . Colocalizes with actin filaments at lamellipodia margins and at the leading edge of migrating cells (PubMed:20052411). In retinal pigment epithelial cells, predominantly localized to stress fiber-like structures with some localization to cytoplasmic puncta (PubMed:27331610). .
<b>Tissue Specificity</b>	In the kidney, expressed in the glomeruli. Also expressed in leukocytes.
<b>Function</b>	
<b>Background</b>	
<b>matters needing attention</b>	Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**