

## IL-13R a 2 Monoclonal Antibody

| Catalog No         | YP-mAb-13371   |
|--------------------|--|
| Isotype            | IgG  |
| Reactivity         | Human;Rat;Mouse;   |
| Applications       | WB   |
| Gene Name          | IL13RA2  |
| Protein Name       | Interleukin-13 receptor subunit alpha-2  |
| Immunogen          | The antiserum was produced against synthesized peptide derived from human IL13RA2. AA range:315-364  |
| Specificity        | IL-13R $\alpha$ 2 Monoclonal Antibody detects endogenous levels of IL-13R $\alpha$ 2 protein.  |
| Formulation        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| Source             | Monoclonal, Mouse,lgG  |
| Purification       | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   |
| Dilution           | WB 1:500-1:2000  |
| Concentration      | 1 mg/ml  |
| Purity             | ≥90%   |
| Storage Stability  | -20°C/1 year   |
| Synonyms           | IL13RA2; IL13R; Interleukin-13 receptor subunit alpha-2; IL-13 receptor subunit alpha-2; IL-13R subunit alpha-2; IL-13R-alpha-2; IL-13RA2; Interleukin-13-binding protein; CD antigen CD213a2  |
| Observed Band      | 50kD   |
| Cell Pathway       | Membrane; Single-pass type I membrane protein.   |
| Tissue Specificity | Brain,Lung,Prostate,Renal cell carcinoma,Testis,   |
| Function           | domain:The box 1 motif is required for JAK interaction and/or activation.,domain:The WSXWS motif appears to be necessary for proper protein folding and thereby efficient intracellular transport and cell-surface receptor binding.,function:Binds as a monomer with high affinity to interleukin 13 (IL13), but not to IL4.,similarity:Belongs to the type I cytokine receptor family. Type 5 subfamily.,similarity:Contains 1 fibronectin type-III domain., |
| Background         | The protein encoded by this gene is closely related to II13RA1, a subuint of the interleukin 13 receptor complex. This protein binds IL13 with high affinity, but lacks cytoplasmic domain, and does not appear to function as a signal mediator. It is reported to play a role in the internalization of IL13. [provided by RefSeq, Jul 2008],  |



## UpingBio technology Co.,Ltd





matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**