







## Olfactory receptor 52N1 Monoclonal Antibody

Background  Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated		
Reactivity Human;Rat;Mouse;  Applications WB  Gene Name OR52N1  Protein Name Olfactory receptor 52N1  Immunogen The antiserum was produced against synthesized peptide derived from human OR52N1. AA range:269-318  Specificity Olfactory receptor 52N1 Monoclonal Antibody detects endogenous levels of Olfactory receptor 52N1 Monoclonal Antibody detects endogenous levels of Olfactory receptor 52N1 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse, lgG  Purification The antibody was affinitly-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms OR52N1; Olfactory receptor 52N1; Olfactory receptor OR11-61  Observed Band 36kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function function: Odorant receptor ., similarity: Belongs to the G-protein coupled receptor family., Background  Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled recoptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many heurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediate the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of oth	Catalog No	YP-mAb-13573
Applications WB  Gene Name OR52N1  Protein Name Olfactory receptor 52N1  Immunogen The antiserum was produced against synthesized peptide derived from human OR52N1. AA range:269-318  Specificity Olfactory receptor 52N1 Monoclonal Antibody detects endogenous levels of Olfactory receptor 52N1 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Diliution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms OR52N1; Olfactory receptor 52N1; Olfactory receptor OR11-61  Observed Band 36kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family.,  Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor protein-coupled receptors arising from single coding-exon genes. Olfactory receptors share a remembers of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a remembers of a large family of G-protein-coupled receptors for this organisms in independent of other organisms. [provided by provided	Isotype	IgG
Gene Name         OR52N1           Protein Name         Olfactory receptor 52N1           Immunogen         The antiserum was produced against synthesized peptide derived from human OR52N1. AA range:269-318           Specificity         Olfactory receptor 52N1 Monoclonal Antibody detects endogenous levels of Olfactory receptor 52N1 protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-1:2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         OR52N1; Olfactory receptor 52N1; Olfactory receptor OR11-61           Observed Band         36kD           Cell Pathway         Cell membrane; Multi-pass membrane protein.           Tissue Specificity         Function           Background         Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated trans	Reactivity	Human;Rat;Mouse;
Protein Name         Olfactory receptor 52N1           Immunogen         The antiserum was produced against synthesized peptide derived from human OR52N1. AA range:269-318           Specificity         Olfactory receptor 52N1 Monoclonal Antibody detects endogenous levels of Olfactory receptor 52N1 protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-1:2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         OR52N1; Olfactory receptor 52N1; Olfactory receptor OR11-61           Observed Band         36kD           Cell Pathway         Cell membrane; Multi-pass membrane protein.           Tissue Specificity         Function           Function         function:Odorant receptor .,simillarity:Belongs to the G-protein coupled receptor family.           Background         Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are	Applications	WB
Immunogen         The antiserum was produced against synthesized peptide derived from human OR52N1. AA range:269-318           Specificity         Olfactory receptor 52N1 Monoclonal Antibody detects endogenous levels of Olfactory receptor 52N1 protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-1:2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         OR52N1; Olfactory receptor 52N1; Olfactory receptor OR11-61           Observed Band         36kD           Cell Pathway         Cell membrane; Multi-pass membrane protein.           Tissue Specificity         Function           Function         function:Odorant receptor .,simillarity:Belongs to the G-protein coupled receptor family.,           Background         Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are a neuronal response that triggers the perception of a smell. The olfactory receptors in a neuronal response that triggers the perception of a protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptor gene family	Gene Name	OR52N1
OR52N1. AA range:269-318  Specificity Olfactory receptor 52N1 Monoclonal Antibody detects endogenous levels of Olfactory receptor 52N1 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90% Storage Stability -20°C/1 year  Synonyms OR52N1; Olfactory receptor 52N1; Olfactory receptor OR11-61  Observed Band 36kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family.,  Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors (GPCR) arising from single coding-exon genes. Olfactory receptor gene family is the largest the genome. The nomenclature assigned to the olfactory receptor gene family is the largest the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Protein Name	Olfactory receptor 52N1
Olfactory receptor 52N1 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms OR52N1; Olfactory receptor 52N1; Olfactory receptor OR11-61  Observed Band 36kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., arising from single coding-exon genes. Olfactory receptor share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant single. The olfactory receptor gene family is the largest the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Immunogen	
Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-1:2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         OR52N1; Olfactory receptor 52N1; Olfactory receptor OR11-61           Observed Band         36kD           Cell Pathway         Cell membrane; Multi-pass membrane protein.           Tissue Specificity         Function           Function         function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family.,           Background         Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are membres of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors (GPCR) arising from single coding-exon genes. Olfactory receptors pare and protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Specificity	Olfactory receptor 52N1 Monoclonal Antibody detects endogenous levels of Olfactory receptor 52N1 protein.
Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-1:2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  OR52N1; Olfactory receptor 52N1; Olfactory receptor OR11-61  Observed Band  36kD  Cell Pathway  Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function  function: Odorant receptor .,similarity: Belongs to the G-protein coupled receptor family.,  Background  Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein—coupled receptors arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein—mediated transduction of odorant signals. The olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms OR52N1; Olfactory receptor 52N1; Olfactory receptor OR11-61  Observed Band 36kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family.,  Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Source	Monoclonal, Mouse,lgG
Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms OR52N1; Olfactory receptor 52N1; Olfactory receptor OR11-61  Observed Band 36kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family.,  Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Purification	
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms OR52N1; Olfactory receptor 52N1; Olfactory receptor OR11-61  Observed Band 36kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family.,  Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Dilution	WB 1:500-1:2000
Synonyms OR52N1; Olfactory receptor 52N1; Olfactory receptor OR11-61  Observed Band 36kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family.,  Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Concentration	1 mg/ml
Synonyms  OR52N1; Olfactory receptor 52N1; Olfactory receptor OR11-61  Observed Band  36kD  Cell Pathway  Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function  function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family.,  Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Purity	≥90%
Observed Band  Cell Pathway  Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function  function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family.,  Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Storage Stability	-20°C/1 year
Cell Pathway  Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function  function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family.,  Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Synonyms	OR52N1; Olfactory receptor 52N1; Olfactory receptor OR11-61
Tissue Specificity  Function  function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family.,  Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Observed Band	36kD
Function  function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family.,  Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Cell Pathway	Cell membrane; Multi-pass membrane protein.
Background  Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Tissue Specificity	
neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Function	function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,
	Background	neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by



## UpingBio technology Co.,Ltd





matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**