



VATL mouse mAb

Catalog No	YP-mAb-17242
Isotype	IgG
Reactivity	Human, Mouse,Rat
Applications	WB
Gene Name	ATP6V0C ATP6C ATP6L ATPL
Protein Name	V-type proton ATPase 16 kDa proteolipid subunit (V-ATPase 16 kDa proteolipid subunit) (Vacuolar proton pump 16 kDa proteolipid subunit)
Immunogen	Synthesized peptide derived from human N-teral VATL
Specificity	This antibody detects endogenous levels of VATL at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Mouse,Monoclonal
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	V-type proton ATPase 16 kDa proteolipid subunit (V-ATPase 16 kDa proteolipid subunit) (Vacuolar proton pump 16 kDa proteolipid subunit)
Observed Band	17kD
Cell Pathway	Cytoplasmic vesicle, clathrin-coated vesicle membrane ; Multi-pass membrane protein . Cytoplasmic vesicle, secretory vesicle, synaptic vesicle membrane ; Multi-pass membrane protein .
Tissue Specificity	
Function	Proton-conducting pore forming of the V0 complex of vacuolar(H ⁺)-ATPase (V-ATPase), a multisubunit enzyme composed of a peripheral complex (V1) that hydrolyzes ATP and a membrane integral complex (V0) that translocates protons . V-ATPase is responsible for acidifying and maintaining the pH of intracellular compartments and in some cell types, is targeted to the plasma membrane, where it is responsible for acidifying the extracellular environment (By similarity).
Background	
matters needing attention	Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images