

KLHL20 mouse mAb

YP-mAb-17918
IgG
Human;Mouse;Rat
WB
KLHL20 KLEIP
Kelch-like protein 20 (Kelch-like ECT2-interacting protein) (Kelch-like protein X)
Synthesized peptide derived from human KLHL20
This antibody detects endogenous levels of KLHL20 at Human, Mouse,Rat
Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Monoclonal, Mouse,IgG
The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
WB 1:500-1:2000
1 mg/ml
≥90%
-20°C/1 year
67kD
Cytoplasm, perinuclear region. Nucleus. Golgi apparatus, trans-Golgi network. Cell projection, axon. Cell projection, dendrite. Localizes in the perinuclear region in normal conditions. Following IFN-alpha or IFN-gamma treatment, it is relocalized and sequestrated to the PML nuclear bodies, preventing DAPK1 ubiquitination (PubMed:20389280).
Substrate-specific adapter of a BCR (BTB-CUL3-RBX1) E3 ubiquitin-protein ligase complex involved in interferon response and anterograde Golgi to endosome transport. The BCR(KLHL20) E3 ubiquitin ligase complex mediates the ubiquitination of DAPK1, leading to its degradation by the proteasome, thereby acting as a negative regulator of apoptosis. The BCR(KLHL20) E3 ubiquitin ligase complex also specifically mediates 'Lys-33'-linked ubiquitination. Involved in anterograde Golgi to endosome transport by mediating 'Lys-33'-linked

Thank you for your recent purchase



UpingBio technology Co.,Ltd





Background

matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images