



DACH1 mouse mAb

Catalog No	YP-mAb-17940
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	DACH1 DACH
Protein Name	Dachshund homolog 1 (Dach1)
Immunogen	Synthesized peptide derived from human DACH1
Specificity	This antibody detects endogenous levels of DACH1 at Human, Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	83kD
Cell Pathway	Nucleus .
Tissue Specificity	Widely expressed. Isoform 2 is found in brain, heart, kidney, liver, leukocytes and spleen. Isoform 3 is found in liver and heart. Isoform 4 is found in spleen.
Function	Transcription factor that is involved in regulation of organogenesis. Seems to be a regulator of SIX1, SIX6 and probably SIX5. Corepression of precursor cell proliferation in myoblasts by SIX1 is switched to coactivation through recruitment of EYA3 to the SIX1-DACH1 complex. Transcriptional activation seems also to involve association of CREBBP. Seems to act as a corepressor of SIX6 in regulating proliferation by directly repressing cyclin-dependent kinase inhibitors, including the p27Kip1 promoter (By similarity). Inhibits TGF-beta signaling through interaction with SMAD4 and NCOR1. Binds to chromatin DNA via its DACHbox-N domain (By similarity).
Background	
matters needing attention	Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images