







## EIF4E2 mouse mAb

Catalog No	YP-mAb-17973
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	EIF4E2 EIF4EL3
Protein Name	Eukaryotic translation initiation factor 4E type 2 (eIF-4E type 2) (eIF4E type 2) (Eukaryotic translation initiation factor 4E homologous protein) (Eukaryotic translation initiation factor 4E-like 3)
Immunogen	Synthesized peptide derived from human EIF4E2
Specificity	This antibody detects endogenous levels of EIF4E2 at Human, Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
o o noon a anon	5
Purity	≥90%
	<u> </u>
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year  27kD
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year  27kD  Cytoplasm . Cytoplasm, P-body .  Recognizes and binds the 7-methylquanosine-containing mRNA can during an
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90%  -20°C/1 year  27kD  Cytoplasm . Cytoplasm, P-body .  Recognizes and binds the 7-methylguanosine-containing mRNA cap during an early step in the initiation . Acts as a repressor of translation initiation . In contrast to EIF4E, it is unable to bind eIF4G (EIF4G1, EIF4G2 or EIF4G3), suggesting that it acts by competing with EIF4E and block assembly of eIF4F at the cap (By similarity). In P-bodies, component of a complex that promotes miRNA-mediated
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90%  -20°C/1 year  27kD  Cytoplasm . Cytoplasm, P-body .  Recognizes and binds the 7-methylguanosine-containing mRNA cap during an early step in the initiation . Acts as a repressor of translation initiation . In contrast to EIF4E, it is unable to bind eIF4G (EIF4G1, EIF4G2 or EIF4G3), suggesting that it acts by competing with EIF4E and block assembly of eIF4F at the cap (By similarity). In P-bodies, component of a complex that promotes miRNA-mediated



## UpingBio technology Co.,Ltd





**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**