





## BTN3A2 mouse mAb

Catalog No	YP-mAb-18004
Isotype	IgG
Reactivity	Human
Applications	WB
Gene Name	BTN3A2 BT3.2 BTF3 BTF4
Protein Name	Butyrophilin subfamily 3 member A2
Immunogen	Synthesized peptide derived from human BTN3A2
Specificity	This antibody detects endogenous levels of BTN3A2 at Human
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
•	
Concentration	1 mg/ml
Purity Purity	1 mg/ml ≥90%
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year  37kD
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year  37kD  Cell membrane ; Single-pass type I membrane protein .
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90%  -20°C/1 year  37kD  Cell membrane ; Single-pass type I membrane protein .  Detected in T-cells and natural killer cells.  Plays a role in T-cell responses in the adaptive immune response. Inhibits the
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90%  -20°C/1 year  37kD  Cell membrane ; Single-pass type I membrane protein .  Detected in T-cells and natural killer cells.  Plays a role in T-cell responses in the adaptive immune response. Inhibits the
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90%  -20°C/1 year  37kD  Cell membrane ; Single-pass type I membrane protein .  Detected in T-cells and natural killer cells.  Plays a role in T-cell responses in the adaptive immune response. Inhibits the release of IFNG from activated T-cells.



## UpingBio technology Co.,Ltd





## **Products Images**