







POLR2L mouse mAb

| Catalog No | YP-mAb-18013 |
|---------------------------|---|
| Isotype | IgG |
| Reactivity | Human;Mouse |
| Applications | WB |
| Gene Name | POLR2L |
| Protein Name | DNA-directed RNA polymerases I, II, and III subunit RPABC5 (RNA polymerases I, II, and III subunit ABC5) (DNA-directed RNA polymerase III subunit L) (RNA polymerase II 7.6 kDa subunit) (RPB7.6) (RPB10 |
| Immunogen | Synthesized peptide derived from human POLR2L |
| Specificity | This antibody detects endogenous levels of POLR2L at Human, Mouse |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,lgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-1:2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | 7kD |
| Cell Pathway | Nucleus . |
| Tissue Specificity | |
| Function | DNA-dependent RNA polymerase catalyzes the transcription of DNA into RNA using the four ribonucleoside triphosphates as substrates. Common component of RNA polymerases I, II and III which synthesize ribosomal RNA precursors, mRNA precursors and many functional non-coding RNAs, and a small RNAs, such as 5S rRNA and tRNAs, respectively. Pol II is the central component of the basal RNA polymerase II transcription machinery. Pols are composed of mobile elements that move relative to each other. In Pol II, POLR2L/RBP10 is part of the core element with the central large cleft (By similarity). |
| Background | |
| matters needing attention | Avoid repeated freezing and thawing! |
| | |



UpingBio technology Co.,Ltd





Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images