



HCP1/PCFT mouse mAb

Catalog No	YP-mAb-18088
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	SLC46A1 HCP1 PCFT
Protein Name	Proton-coupled folate transporter (G21) (Heme carrier protein 1) (PCFT/HCP1) (Solute carrier family 46 member 1)
Immunogen	Synthesized peptide derived from human HCP1/PCFT
Specificity	This antibody detects endogenous levels of HCP1/PCFT at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	50kD
Cell Pathway	Cell membrane ; Multi-pass membrane protein . Apical cell membrane ; Multi-pass membrane protein . Basolateral cell membrane ; Multi-pass membrane protein . Endosome membrane ; Multi-pass membrane protein . Cytoplasm . Localizes to the apical membrane of intestinal cells in iron-deficient cells, while it resides in the cytoplasm in iron-replete cells (By similarity). Localizes to the basolateral membrane of choroid plexus (PubMed:19074442). .
Tissue Specificity	Expressed at highest level in the upper half of the small intestine (duodenum and jejunum), expression decreases downwardly in the subsequent quarter and is undetectable in the last quarter (the lowest ileum) (PubMed:17129779, PubMed:19762432). Also expressed in kidney, liver, placenta, spleen, retina and retinal pigment epithelium (PubMed:17129779, PubMed:17335806). Lower levels found in testis (PubMed:17129779). Very low levels in brain, lung, stomach, heart and muscle (PubMed:17129779).
Function	Proton-coupled folate transporter that mediates folate absorption using an H(+) gradient as a driving force . Catalyzes the intestinal absorption of folates at the brush-border membrane of the proximal jejunum, and the transport from blood to cerebrospinal fluid across the choroid plexus . Functions at acidic pH via alternate outward- and inward-open conformation states . Protonation of residues in the outward open state primes the protein for transport . Binding of folate promotes



breaking of salt bridge network and subsequent closure of the extracellular gate, leading to the inward-open state and release of protons and folate . Also able to transport antifolate drugs, such as methotrexate and pemetrexed, which are established treatments for cancer and autoimmune diseases . Involved in FOLR1-mediated endocytosis by serving as a route of export of folates from acidified endosomes . Also

Background

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images