



DHTKD1 mouse mAb

Catalog No	YP-mAb-18159
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	DHTKD1 KIAA1630
Protein Name	Probable 2-oxoglutarate dehydrogenase E1 component DHKTD1, mitochondrial (EC 1.2.4.2) (Dehydrogenase E1 and transketolase domain-containing protein 1)
Immunogen	Synthesized peptide derived from human DHTKD1
Specificity	This antibody detects endogenous levels of DHTKD1 at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	101kD
Cell Pathway	Mitochondrion .
Tissue Specificity	
Function	2-oxoadipate dehydrogenase (E1a) component of the 2-oxoadipate dehydrogenase complex (OADHC) . Participates in the first step, rate limiting for the overall conversion of 2-oxoadipate (alpha-ketoadipate) to glutaryl-CoA and CO(2) catalyzed by the whole OADHC . Catalyzes the irreversible decarboxylation of 2-oxoadipate via the thiamine diphosphate (ThDP) cofactor and subsequent transfer of the decarboxylated acyl intermediate on an oxidized dihydrolipoyl group that is covalently amidated to the E2 enzyme (dihydrolipoyllysine-residue succinyltransferase or DLST) (Probable) . Can catalyze the decarboxylation of 2-oxoglutarate in vitro, but at a much lower rate than 2-oxoadipate . Responsible for the last step of L-lysine, L-hydroxylysine and L-tryptophan catabolism with the common product being 2-oxoadipate (Probable).
Background	

**matters needing attention**

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images