



FLNB mouse mAb

Catalog No	YP-mAb-18271
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	FLNB FLN1L FLN3 TABP TAP
Protein Name	Filamin-B (FLN-B) (ABP-278) (ABP-280 homolog) (Actin-binding-like protein) (Beta-filamin) (Filamin homolog 1) (Fh1) (Filamin-3) (Thyroid autoantigen) (Truncated actin-binding protein) (Truncated ABP)
Immunogen	Synthesized peptide derived from human FLNB
Specificity	This antibody detects endogenous levels of FLNB at Human
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	286kD
Cell Pathway	[Isoform 1]: Cytoplasm, cell cortex. Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, stress fiber. Cytoplasm, myofibril, sarcomere, Z line. In differentiating myotubes, isoform 1, isoform 2 and isoform 3 are localized diffusely throughout the cytoplasm with regions of enrichment at the longitudinal actin stress fiber. In differentiated tubes, isoform 1 is also detected within the Z-lines.; [Isoform 2]: Cytoplasm, cytoskeleton, stress fiber.; [Isoform 3]: Cytoplasm, cytoskeleton, stress fiber.; [Isoform 6]: Cytoplasm, cytoskeleton. Polarized at the periphery of myotubes.
Tissue Specificity	Ubiquitous. Isoform 1 and isoform 2 are expressed in placenta, bone marrow, brain, umbilical vein endothelial cells (HUVEC), retina and skeletal muscle. Isoform 1 is predominantly expressed in prostate, uterus, liver, thyroid, stomach, lymph node, small intestine, spleen, skeletal muscle, kidney, placenta, pancreas, heart, lung, platelets, endothelial cells, megakaryocytic and erythroleukemic cell lines. Isoform 2 is predominantly expressed in spinal cord, platelet and Daudi cells. Also expressed in thyroid adenoma, neurofibrillary tangles (NFT), senile plaques in the hippocampus and cerebral cortex in Alzheimer disease (AD). Isoform 3 and isoform 6 are expressed predominantly in lung, heart, skeletal muscle, testis, spleen, thymus and leukocytes. Isoform 4 and isoform 5 are expressed in h

**Function**

Connects cell membrane constituents to the actin cytoskeleton. May promote orthogonal branching of actin filaments and links actin filaments to membrane glycoproteins. Anchors various transmembrane proteins to the actin cytoskeleton. Interaction with FLNA may allow neuroblast migration from the ventricular zone into the cortical plate. Various interactions and localizations of isoforms affect myotube morphology and myogenesis. Isoform 6 accelerates muscle differentiation in vitro.

Background**matters needing attention**

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images