



# AHCYL1 Rabbit pAb

Catalog No	YP-Ab-18574
Isotype	IgG
Reactivity	Human, Mouse
Applications	WB
Gene Name	AHCYL1 DCAL XPVKONA
Protein Name	Putative adenosylhomocysteinase 2 (AdoHcyase 2) (DC-expressed AHCY-like molecule) (S-adenosyl-L-homocysteine hydrolase 2) (S-adenosylhomocysteine hydrolase-like protein 1)
Immunogen	Synthesized peptide derived from human AHCYL1
Specificity	This antibody detects endogenous levels of AHCYL1 at Human, Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	58kD
Cell Pathway	Endoplasmic reticulum . Cytoplasm, cytosol . Apical cell membrane ; Peripheral membrane protein . Microsome . Associates with membranes when phosphorylated, probably through interaction with ITPR1 (By similarity). Localizes to mitochondria-associated endoplasmic reticulum membranes (MAMs) (PubMed:27995898). Localization to MAMs is greatly reduced under apoptotic stress conditions (PubMed:27995898). .
Tissue Specificity	Expressed in dendritic cells.
Function	Multifaceted cellular regulator which coordinates several essential cellular functions including regulation of epithelial HCO <sub>3</sub> <sup>-</sup> and fluid secretion, mRNA processing and DNA replication. Regulates ITPR1 sensitivity to inositol 1,4,5-trisphosphate, competing for the common binding site and acting as endogenous 'pseudoligand' whose inhibitory activity can be modulated by its phosphorylation status. Promotes the formation of contact points between the endoplasmic reticulum (ER) and mitochondria, facilitating transfer of Ca <sup>2+</sup> from the ER to mitochondria . Under normal cellular conditions, functions cooperatively with BCL2L10 to limit ITPR1-mediated Ca <sup>2+</sup> release but, under apoptotic stress conditions, dephosphorylated which promotes dissociation of both AHCYL1 and



BCL2L10 from mitochondria-associated endoplasmic reticulum membranes, inhibits BCL2L10 interaction with ITPR1 and leads to increased  $\text{Ca}^{2+}$  transfer to mitochondria which promotes apoptosis. In the pancreatic and salivary ducts, at resting state, attenuates inositol 1,4,5-trisphosphate-induced calcium release by interacting with ITPR1. When extracellular stimuli induce ITPR1 phosphorylation or inositol 1,4,5-trisphosphate production, dissociates from ITPR1 to interact with CFTR and SLC26A6, mediating their synergistic activation by calcium and cAMP that stimulates the epithelial secretion of electrolytes and fluid (By similarity). Also activates basolateral SLC4A4 isoform 1 to coordinate fluid and  $\text{HCO}_3^-$  secretion. Inhibits the effect of STK39 on SLC4A4 and CFTR by recruiting PP1 phosphatase which activates SLC4A4, SLC26A6 and CFTR through dephosphorylation (By similarity). Mediates the induction of SLC9A3 surface expression produced by Angiotensin-2. Depending on the cell type, activates SLC9A3 in response to calcium or reverses SLC9A3R2-dependent calcium inhibition. May modulate the polyadenylation state of specific mRNAs, both by controlling the subcellular location of FIP1L1 and by inhibiting PAPOLA activity, in response to a stimulus that

## Background

### matters needing attention

Avoid repeated freezing and thawing!

### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images