

**(** Tel: 400-999-8863 ■ Email:Upingbio.163.com





## Angptl1 Polyclonal Antibody

| Catalog No         | YP-Ab-16025  |
|--------------------|--|
| Isotype            | IgG  |
| Reactivity         | Human;Mouse;Rat  |
| Applications       | WB;IHC;IF;ELISA  |
| Gene Name          | ANGPTL1  |
| Protein Name       | Angiopoietin-related protein 1   |
| Immunogen          | The antiserum was produced against synthesized peptide derived from the Internal region of human ANGPTL1. AA range:11-60   |
| Specificity        | Angptl1 Polyclonal Antibody detects endogenous levels of Angptl1 protein.  |
| Formulation        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| Source             | Polyclonal, Rabbit,IgG   |
| Purification       | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  |
| Dilution           | WB: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000 IF 1:50-200   |
| Concentration      | 1 mg/ml  |
| Purity             | ≥90%   |
| Storage Stability  | -20°C/1 year   |
| Synonyms           | ANGPTL1; ANG3; ANGPT3; ARP1; PSEC0154; Angiopoietin-related protein 1; Angiopoietin-3; ANG-3; Angiopoietin-like protein 1  |
| Observed Band      | 57kD   |
| Cell Pathway       | Secreted .   |
| Tissue Specificity | Highly expressed in adrenal gland, placenta, thyroid gland, heart, skeletal muscle and small intestine. Weakly expressed in testis, ovary, colon, pancreas, kidney and stomach.  |
| Function           | similarity:Contains 1 fibrinogen C-terminal domain.,tissue specificity:Highly expressed in adrenal gland, placenta, thyroid gland, heart, skeletal muscle and small intestine. Weakly expressed in testis, ovary, colon, pancreas, kidney and stomach.,  |
| Background         | Angiopoietins are members of the vascular endothelial growth factor family and the only known growth factors largely specific for vascular endothelium. Angiopoietin-1, angiopoietin-2, and angiopoietin-4 participate in the formation of blood vessels. The protein encoded by this gene is another member of the angiopoietin family that is widely expressed in adult tissues with mRNA levels highest in highly vascularized tissues. This protein was found to be a secretory protein that does not act as an endothelial cell mitogen in vitro. [provided by RefSeq, Jul 2008], |



## UpingBio technology Co.,Ltd

C Tel: 400-999-8863 🗷 Email:Upingbio.163.com




matters needing attention

Avoid repeated freezing and thawing!


**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


## **Products Images**



Western Blot analysis of K562 cells using Angptl1 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded mouse-heart, antibody was diluted at 1:100




Immunohistochemical analysis of paraffin-embedded mouse-heart, antibody was diluted at 1:100



## UpingBio technology Co.,Ltd

**(** Tel: 400-999-8863 **(** Emall:Upingbio.163.com





Western blot analysis of lysate from K562 cells, using ANGPTL1 Antibody.