



AGS3 Polyclonal Antibody

Catalog No	YP-Ab-16126
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	IHC;IF;ELISA
Gene Name	GPSM1
Protein Name	G-protein-signaling modulator 1
Immunogen	Synthesized peptide derived from the Internal region of human AGS3.
Specificity	AGS3 Polyclonal Antibody detects endogenous levels of AGS3 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	IHC: 1/100 - 1/300. ELISA: 1/5000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	GPSM1; AGS3; G-protein-signaling modulator 1; Activator of G-protein signaling 3
Observed Band	
Cell Pathway	Cytoplasm, cytosol. Endoplasmic reticulum membrane; Peripheral membrane protein; Cytoplasmic side. Golgi apparatus membrane; Peripheral membrane protein; Cytoplasmic side. Cell membrane; Peripheral membrane protein; Cytoplasmic side.
Tissue Specificity	Expressed in intestinal cells.
Function	domain:The GoLoco domains mediate interaction with G(i/o) alpha (By similarity). The GoLoco domains are essential for the GDI activity toward G(i/o) alpha.,function:Guanine nucleotide dissociation inhibitor (GDI) which functions as a receptor-independent activator of heterotrimeric G-protein signaling. Keeps G(i/o) alpha subunit in its GDP-bound form thus uncoupling heterotrimeric G-proteins signaling from G protein-coupled receptors. Controls spindle orientation and asymmetric cell fate of cerebral cortical progenitors. May also be involved in macroautophagy in intestinal cells. May play a role in drug addiction.,PTM:Phosphorylation regulates interaction with G(i/o) alpha.,similarity:Belongs to the GPSM family.,similarity:Contains 4 GoLoco domains.,similarity:Contains 9 TPR repeats.,subunit:Interacts with GNAI1, GNAI2 and GNAI3 preferentially in their GDP-bound state. May also interact



Background

G-protein signaling modulators (GPSMs) play diverse functional roles through their interaction with G-protein subunits. This gene encodes a receptor-independent activator of G protein signaling, which is one of several factors that influence the basal activity of G-protein signaling systems. The protein contains seven tetratricopeptide repeats in its N-terminal half and four G-protein regulatory (GPR) motifs in its C-terminal half. Multiple alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2011],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded human Colon cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).