

(Tel: 400-999-8863 ■ Emall:Upingbio.163.com



Calnexin Monoclonal Antibody

Catalog No	YP-Ab-16267
Isotype	IgG
Reactivity	Human
Applications	WB;IF;ELISA
Gene Name	CANX
Protein Name	Calnexin
Immunogen	Synthetic peptide corresponding to aa (CEAAEERPWLWVVYILTVAL) of human Calnexin, conjugated to KLH.
Specificity	Calnexin Monoclonal Antibody detects endogenous levels of Calnexin protein.
Formulation	Ascitic fluid containing 0.03% sodium azide,0.5% BSA, 50%glycerol.
Source	Monoclonal, Mouse
Purification	Affinity purification
Dilution	Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Purity Storage Stability	≥90% -20°C/1 year
Storage Stability	-20°C/1 year CANX; Calnexin; IP90; Major histocompatibility complex class I antigen-binding
Storage Stability Synonyms	-20°C/1 year CANX; Calnexin; IP90; Major histocompatibility complex class I antigen-binding
Storage Stability Synonyms Observed Band	-20°C/1 year CANX; Calnexin; IP90; Major histocompatibility complex class I antigen-binding protein p88; p90 Endoplasmic reticulum membrane; Single-pass type I membrane protein. Endoplasmic reticulum. Melanosome. Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:12643545, PubMed:17081065). The palmitoylated form preferentially localizes to the
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year CANX; Calnexin; IP90; Major histocompatibility complex class I antigen-binding protein p88; p90 Endoplasmic reticulum membrane; Single-pass type I membrane protein. Endoplasmic reticulum. Melanosome. Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:12643545, PubMed:17081065). The palmitoylated form preferentially localizes to the perinuclear rough ER (PubMed:22314232). B-cell lymphoma, Epithelium, Fibroblast, Keratinocyte, Kidney, Liver, Lymph, Placenta, Plate



UpingBio technology Co.,Ltd

C Tel: 400-999-8863 🗷 Email:Upingbio.163.com



facilitating protein folding and assembly. It may also play a central role in the quality control of protein folding by retaining incorrectly folded protein subunits within the ER for degradation. Alternatively spliced transcript variants encoding the same protein have been described. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis using Calnexin Monoclonal Antibody against A431 (1), HeLa (2), MCF-7 (3) and A549 (4) cell lysate.



Confocal immunofluorescence analysis of Hela cells using Calnexin Monoclonal Antibody (green).