



# Creatine Kinase M Polyclonal Antibody

<b>Catalog No</b>	YP-Ab-16716
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	WB;IHC;IF;ELISA
<b>Gene Name</b>	CKM
<b>Protein Name</b>	Creatine kinase M-type
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human M-CK. AA range:10-59
<b>Specificity</b>	Creatine Kinase M Polyclonal Antibody detects endogenous levels of Creatine Kinase M protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	CKM; CKMM; Creatine kinase M-type; Creatine kinase M chain; M-CK
<b>Observed Band</b>	43kD
<b>Cell Pathway</b>	Cytoplasm.
<b>Tissue Specificity</b>	Liver,
<b>Function</b>	catalytic activity:ATP + creatine = ADP + phosphocreatine.,function:Reversibly catalyzes the transfer of phosphate between ATP and various phosphogens (e.g. creatine phosphate). Creatine kinase isoenzymes play a central role in energy transduction in tissues with large, fluctuating energy demands, such as skeletal muscle, heart, brain and spermatozoa.,online information:CKM entry,online information:Creatine kinase entry,similarity:Belongs to the ATP:guanido phosphotransferase family.,subunit:Dimer of identical or non-identical chains. With MM being the major form in skeletal muscle and myocardium, MB existing in myocardium, and BB existing in many tissues, especially brain.,
<b>Background</b>	The protein encoded by this gene is a cytoplasmic enzyme involved in energy homeostasis and is an important serum marker for myocardial infarction. The encoded protein reversibly catalyzes the transfer of phosphate between ATP and various phosphogens such as creatine phosphate. It acts as a homodimer in



striated muscle as well as in other tissues, and as a heterodimer with a similar brain isozyme in heart. The encoded protein is a member of the ATP:guanido phosphotransferase protein family. [provided by RefSeq, Jul 2008],


#### matters needing attention

Avoid repeated freezing and thawing!


#### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


### Products Images




Western Blot analysis of various cells using Creatine Kinase M Polyclonal Antibody




Western Blot analysis of Jurkat cells using Creatine Kinase M Polyclonal Antibody



The picture was kindly provided by our customer



Immunofluorescence analysis of HepG2 cells, using M-CK Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human heart tissue, using M-CK Antibody. The picture on the right is blocked with the synthesized peptide.